

Topological insulators from the perspective of non-commutative geometry and index theory

Hermann Schulz-Baldes
Erlangen

main collaborators:

Prodan, Loring, Carey, Grossmann, Phillips
De Nittis, Villegas, Kellendonk, Richter, Bellissard

Graduiertenkolleg 1838
Freudenstadt
February 2017

Plan for the lectures

- What is a topological insulator?
 - What are the main experimental facts?
 - What are the main theoretical elements?
 - Almost everything in a one-dimensional toy model (SSH model)
 - Toy models for higher dimension
 - Algebraic formalism (crossed product C^* -algebras)
 - Measurable quantities as topological invariants
 - Bulk-edge correspondence
 - Index theorems for invariants
 - Implementation of symmetries (periodic table of topological ins.)
- Math tools:** K -theory, index theory and non-commutative geometry

1. Experimental facts
2. Elements of basic theory
3. One-dimensional toy model
4. K -theory crash course
5. Observable algebra for tight-binding models
6. Topological invariants in solid state systems
7. Invariants as response coefficients
8. Bulk-boundary correspondence
9. Implementation of symmetries
10. Laughlin arguments
11. Dirty superconductors

1 Experimental facts

What is a topological insulator?

- d -dimensional disordered system of independent Fermions with a combination of basic symmetries


TRS, PHS, CHS = time reversal, particle hole, chiral symmetry

- Fermi level in a Gap or Anderson localization regime
- Topology of bulk (in Bloch bundles over Brillouin torus):



winding numbers, Chern numbers, \mathbb{Z}_2 -invariants, higher invariants

- Delocalized edge modes with non-trivial topology
- Bulk-edge correspondence
- Topological bound states at defects (zero modes)
- Toy models: tight-binding Hamiltonians
- Wider notions include interactions, bosons, spins, photonic crys.


Quantum Hall Effect: first topological insulator



Schematic representation of IQHE



Most important facts for IQHE



Two-dimensional electron gas between two doped semiconductors (Spot error in picture!) Measure of macroscopic (!) Hall tension

$$\sigma = \frac{I_{x,x}}{V_{x,y}} = n \frac{e^2}{h} \quad \text{with } n \in \mathbb{N}$$

Integer quantization with relative error 10^{-8} with fundamental constant

Strong magnetic field and electron density can be modified

Anderson localized states can be filled without changing conductivity

Prizes and further advances on the QHE

Nobel prizes:

- Klitzing (1985)
- Störmer-Tsui-Laughlin (1998) for fractional QHE
- Thouless (2016) explanation of integer QHE & Thouless-Kosterlitz
- Haldane (2016) anomalous QHE & Haldane spin chain
NO exterior magnetic field, only magnetic material
- QHE in graphene at room temperature
Novoselov, Geim et al 2007 (Nobel 2005)
- Anomalous QHE at room temperature in SnGe (Chinese group 2016)
Review: Ren, Qiao, Niu 2016

Quantum spin Hall systems

Prior to 2005: no (local) magnetic field, no topology

Kane-Mele (2005):

\mathbb{Z}_2 -topology in two-dimensional systems with time-reversal symmetry

First erroneous proposal: spin orbit coupling in graphene (too small)

Theoretical prediction by Bernevig and Zhang (2006): look into HgTe

Measurement by Molenkamp group in Würzburg

Complicated samples, inconsistencies with theory, so still disputed

Measurement in more conventional Si-semiconductor by Du group
(Rice 2014) Surprise: stability w.r.t. magnetic field

Majorana zero modes

First proposal (Read-Green 2000):

attached to flux tubes in 2d ($p + ip$)-wave superconductors

Second proposal (Kitaev, Beenacker group, Alicea, *etc.*):

at ends of dirty superconductor wires placed on a semiconductor

Measurement in C. Marcus group (2014-2016 Bohr Inst., Kopenhagen)

Further measurements in Delft and Princeton groups

2017: <http://www.seethroughthe.cloud/2017/01/23/>

Headline is: Microsoft Steps Away From The Chalk Board
to Create Quantum Computer

Mysterious citation:

*The magic recipe involves a combination of
semiconductors and superconductors*

Higher dimensional topological insulators?

Table I. Summary of topological insulator materials that have been experimentally addressed. The definition of (1;111) etc. is introduced in Sect. 3.7. (In this table, S.S., P.T., and SM stand for surface state, phase transition, and semimetal, respectively.)

Type	Material	Band gap	Bulk transport	Remark	Reference
2D, $\nu = 1$	CdTe/HgTe/CdTe	<10 meV	insulating	high mobility	31
2D, $\nu = 1$	AlSb/InAs/GaSb/AlSb	~4 meV	weakly insulating	gap is too small	73
3D (1;111)	$\text{Bi}_{1-x}\text{Sb}_x$	<30 meV	weakly insulating	complex S.S.	36, 40
3D (1;111)	Sb	semimetal	metallic	complex S.S.	39
3D (1;000)	Bi_2Se_3	0.3 eV	metallic	simple S.S.	94
3D (1;000)	Bi_2Te_3	0.17 eV	metallic	distorted S.S.	95, 96
3D (1;000)	Sb_2Te_3	0.3 eV	metallic	heavily p-type	97
3D (1;000)	$\text{Bi}_2\text{Te}_2\text{Se}$	~0.2 eV	reasonably insulating	ρ_{xx} up to 6 Ω cm	102, 103, 105
3D (1;000)	$(\text{Bi,Sb})_2\text{Te}_3$	<0.2 eV	moderately insulating	mostly thin films	193
3D (1;000)	$\text{Bi}_{2-x}\text{Sb}_x\text{Te}_{3-y}\text{Se}_y$	<0.3 eV	reasonably insulating	Dirac-cone engineering	107, 108, 212
3D (1;000)	$\text{Bi}_2\text{Te}_{1.6}\text{S}_{1.4}$	0.2 eV	metallic	n-type	210
3D (1;000)	$\text{Bi}_{1.1}\text{Sb}_{0.9}\text{Te}_2\text{S}$	0.2 eV	moderately insulating	ρ_{xx} up to 0.1 Ω cm	210
3D (1;000)	$\text{Sb}_2\text{Te}_2\text{Se}$?	metallic	heavily p-type	102
3D (1;000)	$\text{Bi}_2(\text{Te,Se})_2(\text{Se,S})$	0.3 eV	semi-metallic	natural Kawazulite	211
3D (1;000)	TlBiSe ₂	~0.35 eV	metallic	simple S.S., large gap	110–112
3D (1;000)	TlBiTe ₂	~0.2 eV	metallic	distorted S.S.	112
3D (1;000)	TlBi(S,Se) ₂	<0.35 eV	metallic	topological P.T.	116, 117
3D (1;000)	PbBi_2Te_4	~0.2 eV	metallic	S.S. nearly parabolic	121, 124
3D (1;000)	PbSb_2Te_4	?	metallic	p-type	121
3D (1;000)	GeBi_2Te_4	0.18 eV	metallic	n-type	102, 119, 120
3D (1;000)	PbBi_4Te_7	0.2 eV	metallic	heavily n-type	125
3D (1;000)	$\text{GeBi}_{4-x}\text{Sb}_x\text{Te}_7$	0.1–0.2 eV	metallic	n (p) type at $x = 0$ (1)	126
3D (1;000)	$(\text{PbSe})_5(\text{Bi}_2\text{Se}_3)_6$	0.5 eV	metallic	natural heterostructure	130
3D (1;000)	$(\text{Bi}_2)_n(\text{Bi}_2\text{Se}_3)_m$	semimetal	metallic	$(\text{Bi}_2)_n(\text{Bi}_2\text{Se}_3)_m$ series	127

2 Elements of basic theory

First for QHE in continuous physical space:

Landau-operator with disordered potential

$$H = \frac{1}{2m}(i\partial_{x_1} - eA_1)^2 + \frac{1}{2m}(i\partial_{x_2} - eA_2)^2 + \lambda V_{\text{dis}}$$

on Hilbert space $L^2(\mathbb{R}^2)$. Landau gauge $A_1 = 0$ and $A_2 = BX_1$

If there is no disorder $\lambda = 0$, Fourier transform in 2-direction works

$$\mathcal{F}_2 H \mathcal{F}_2^* = \int_{\mathbb{R}}^{\oplus} dk_2 H(k_2)$$

with $H(k_2) = H(k_2)^*$ shifted one-dimensional harmonic oscillator

\implies infinitely degenerate so-called Landau bands.

Projection P on lowest band has integral kernel with Hall conductance

$$\begin{aligned} \text{Ch}(P) &= 2\pi i \langle 0 | P[i[X_1, P], i[X_2, P]] | 0 \rangle \\ &= \pi \int_{\mathbb{C}} dx \int_{\mathbb{C}} dy e^{-\frac{1}{2}(|x|^2 + |y|^2 - x\bar{y})} (x\bar{y} - y\bar{x}) = -1 \end{aligned}$$

Effect of disorder


Typical model from i.i.d. $\omega_n \in [-1, 1]$ and $v \in \mathcal{D}$ with $\|v\|_\infty \leq 1$

$$V_{\text{dis}}(x) = \sum_{n \in \mathbb{Z}^2} \omega_n v(x - n)$$

Landau band widens by $\lambda \neq 0$. Gap closes at $\lambda \approx 1$

Expectation: all states Anderson localized, except at one energy

Proof at band edges by Barbaroux, Combes, Hislop 1997, others...



Spectrum of edge states



\hat{H}_L half-space restriction on $L^2(\mathbb{R}_{\geq 0} \times \mathbb{R})$ with Dirichlet

Still without disorder, Fourier transform works also for half-space:

$$\mathcal{F}_2 \hat{H} \mathcal{F}_2^* = \int_{\mathbb{R}}^{\oplus} dk_2 \hat{H}(k_2)$$

with $\hat{H}(k_2) = \hat{H}(k_2)^*$ cut off shifted harmonic oscillator on $L^2(\mathbb{R}_{\geq 0})$

Read off basic bulk-edge correspondence (right pic for generic gap)



Harper model

This is a lattice or tight-binding model on $\ell^2(\mathbb{Z}^2)$


$$H = U_1 + U_1^* + U_2 + U_2^*$$

Here $U_1 = S_1$ shift in 1-direction, and $U_2 = e^{iBX_1} S_2$ (Landau gauge)

Plotted: spectrum as a function of B (Hofstadter's butterfly)

Spectrum fractal for irrational B . Most gaps close with V_{dis}

In each gap there are edge state bands (on $\ell^2(\mathbb{Z} \times \mathbb{N})$, Hatsugai 1993)




Coloured Hofstadter butterfly (Avron, Osadchy)

For each Fermi energy μ one has $P = \chi(H \leq \mu)$

If μ in gap, then Chern number well-defined

$$\text{Ch}(P) = 2\pi i \langle 0 | P[i[X_1, P], i[X_2, P]] | 0 \rangle \in \mathbb{Z}$$

Different values, different colours




Haldane model for anomalous QHE

On honeycomb lattice = decorated triangular lattice, so on $\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^2$

$$H_{\text{Hal}} = M \begin{pmatrix} 0 & S_1 + S_2 + 1 \\ S_1^* + S_2^* + 1 & 0 \end{pmatrix} + t_2 \sum_{j=1}^3 \begin{pmatrix} e^{i\phi_j} S_j + (e^{i\phi_j} S_j)^* & 0 \\ 0 & e^{-i\phi_j} S_j + (e^{-i\phi_j} S_j)^* \end{pmatrix}$$

Here $S_3 = S_1^* S_2$ and $\phi_1 = -\phi_2 = \phi_3 = \phi$. Only periodic magnetic field

Then central gap with $P = \chi(H \leq 0)$ and Chern number $C_1 = \text{Ch}(P)$



Kane-Mele model for SQHE

On honeycomb lattice with spin $\frac{1}{2}$, so on $\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^4$

$$H_{\text{KM}} = \begin{pmatrix} H_{\text{Hal}} & 0 \\ 0 & H_{\text{Hal}} \end{pmatrix} + H_{\text{Ras}}$$


First term comes from spin-orbit coupling to next nearest neighbors

Second Rashba spin-orbit term is off-diagonal breaks chiral symmetry

If H_{Ras} small, central gap still open

Chern number vanishes (TRS), but non-trivial \mathbb{Z}_2 -invariant

This leads to edge states



Discrete symmetries (invoking real structure)

Given commuting real, skew- or selfadjoint unitaries J_{ch} , S_{tr} , S_{ph}

$$\text{chiral symmetry (CHS)} : \quad J_{\text{ch}}^* H J_{\text{ch}} = -H$$

$$\text{time reversal symmetry (TRS)} : \quad S_{\text{tr}}^* \bar{H} S_{\text{tr}} = H$$

$$\text{particle-hole symmetry (PHS)} : \quad S_{\text{ph}}^* \bar{H} S_{\text{ph}} = -H$$

$S_{\text{tr}} = e^{i\pi s^y}$ orthogonal on \mathbb{C}^{2s+1} with $S_{\text{tr}}^2 = \pm 1$ even or odd

S_{ph} orthogonal on \mathbb{C}_{ph}^2 with $S_{\text{ph}}^2 = \pm 1$ even or odd

So typical Hamiltonian acts on $\ell^2(\mathbb{Z}^d) \otimes \mathbb{C}^N \otimes \mathbb{C}^{2s+1} \otimes \mathbb{C}_{\text{ph}}^2$

Note: TRS + PHS \implies CHS with $J_{\text{ch}} = S_{\text{tr}} S_{\text{ph}}$

10 combinations of symmetries: none (1), one (5), three (4)

10 Cartan-Altland-Zirnbauer classes (CAZ): 2 complex, 8 real

Further distinction in each of the 10 classes: topological insulators

Periodic table of topological insulators

Schnyder-Ryu-Furusaki-Ludwig, Kitaev 2008: just strong invariants

$j \setminus d$	TRS	PHS	CHS	1	2	3	4	5	6	7	8
0	0	0	0		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}
1	0	0	1	\mathbb{Z}		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}	
0	+1	0	0				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}
1	+1	+1	1	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2
2	0	+1	0	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2
3	-1	+1	1	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$	
4	-1	0	0		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$
5	-1	-1	1	$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}			
6	0	-1	0		$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}		
7	+1	-1	1			$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}	

3 One-dimensional toy model (SSH, see [PS])

Su-Schrieffer-Heeger (1980, conducting polyacetylene polymer)

$$H = \frac{1}{2}(\sigma_1 + i\sigma_2) \otimes S + \frac{1}{2}(\sigma_1 - i\sigma_2) \otimes S^* + m\sigma_2 \otimes \mathbf{1}$$

where S bilateral shift on $\ell^2(\mathbb{Z})$, $m \in \mathbb{R}$ mass and Pauli matrices

In their grading

$$H = \begin{pmatrix} 0 & S - im \\ S^* + im & 0 \end{pmatrix} \quad \text{on } \ell^2(\mathbb{Z}) \otimes \mathbb{C}^2$$

Off-diagonal \cong chiral symmetry $\sigma_3^* H \sigma_3 = -H$. In Fourier space:

$$H = \int^{\oplus} dk H_k \quad H_k = \begin{pmatrix} 0 & e^{-ik} - im \\ e^{ik} + im & 0 \end{pmatrix}$$

Topological invariant for $m \neq -1, 1$

$$\text{Wind}(k \mapsto e^{ik} + im) = \delta(m \in (-1, 1))$$

Chiral bound states

Half-space Hamiltonian

$$\hat{H} = \begin{pmatrix} 0 & \hat{S} - im \\ \hat{S}^* + im & 0 \end{pmatrix} \quad \text{on } \ell^2(\mathbb{N}) \otimes \mathbb{C}^2$$

where \hat{S} unilateral right shift on $\ell^2(\mathbb{N})$

Still chiral symmetry $\sigma_3^* \hat{H} \sigma_3 = -\hat{H}$

If $m = 0$, simple bound state at $E = 0$ with eigenvector $\psi_0 = \begin{pmatrix} |0\rangle \\ 0 \end{pmatrix}$.

Perturbations, e.g. in m , cannot move or lift this bound state $\psi_m!$

Positive chirality conserved: $\sigma_3 \psi_m = \psi_m$

Theorem 3.1 (Basic bulk-boundary correspondence)

If \hat{P} projection on bound states of \hat{H} , then

$$\text{Wind}(k \mapsto e^{ik} + im) = \text{Tr}(\hat{P}\sigma_3)$$

Disordered model

Add i.i.d. random mass term $\omega = (m_n)_{n \in \mathbb{Z}}$:

$$H_\omega = H + \sum_{n \in \mathbb{Z}} m_n \sigma_2 \otimes |n\rangle\langle n|$$

Still chiral symmetry $\sigma_3^* H_\omega \sigma_3 = -H_\omega$ so

$$H_\omega = \begin{pmatrix} 0 & A_\omega^* \\ A_\omega & 0 \end{pmatrix}$$

Bulk gap at $E = 0 \implies A_\omega$ invertible

Non-commutative winding number, also called first Chern number:

$$\text{Wind} = \text{Ch}_1(A) = i \mathbf{E}_\omega \text{Tr} \langle 0 | A_\omega^{-1} i [X, A_\omega] | 0 \rangle$$

where \mathbf{E}_ω is average over probability measure \mathbb{P} on i.i.d. masses

Index theorem and bulk-boundary correspondence

Theorem 3.2 (Disordered Noether-Gohberg-Krein Theorem)

If Π is Hardy projection on positive half-space, then \mathbb{P} -almost surely

$$\text{Wind} = \text{Ch}_1(A) = -\text{Ind}(\Pi A_\omega \Pi)$$

For periodic model as above, $A_\omega = e^{ik} \in C(\mathbb{S}^1)$

Fredholm operator is then standard Toeplitz operator

Theorem 3.3 (Disordered bulk-boundary correspondence)

If \hat{P}_ω projection on bound states of \hat{H}_ω , then

$$\text{Wind} = \text{Ch}_1(A) = \text{Ch}_0(\hat{P}_\omega) = \text{Tr}(\hat{P}_\omega \sigma_3)$$

Structural robust result:

holds for chiral Hamiltonians with larger fiber, other disorder, etc.

Index in linear algebra

Rank theorem for $T \in \text{Mat}(N \times M, \mathbb{C})$

$$\begin{aligned}M &= \dim(\text{Ker}(T)) + \dim(\text{Ran}(T)) \\ &= \dim(\text{Ker}(T)) + \dim(\text{Ker}(T^*)^\perp) \\ &= \dim(\text{Ker}(T)) + (N - \dim(\text{Ker}(T^*)))\end{aligned}$$

Hence stability of index defined by

$$\text{Ind}(T) = \dim(\text{Ker}(T)) - \dim(\text{Ker}(T^*)) = M - N$$

Homotopy invariance: under continuous perturbation $t \in \mathbb{R} \mapsto T_t$

$$t \in \mathbb{R} \mapsto \text{Ind}(T_t) \text{ konstant}$$

For quadratic matrices, *i.e.* $N = M$, always $\text{Ind}(T) = 0$

Index in infinite dimension

Definition 3.4

$T \in \mathcal{B}(\mathcal{H})$ continuous Fredholm operator on \mathcal{H}

$\iff T\mathcal{H}$ closed, $\dim(\text{Ker}(T)) < \infty$, $\dim(\text{Ker}(T^*)) < \infty$

Then: $\text{Ind}(T) = \dim(\text{Ker}(T)) - \dim(\text{Ker}(T^*))$

Theorem 3.5 (Dieudonné, Krein)

Ind is a compactly stable homotopy invariant:

$$\text{Ind}(T) = \text{Ind}(T + K) = \text{Ind}(T_t)$$

Example: shift $\hat{S} : \ell^2(\mathbb{N}) \rightarrow \ell^2(\mathbb{N})$ by $\hat{S}\psi = (\psi_{n-1})_{n \in \mathbb{N}}$ on $\psi = (\psi_n)_{n \in \mathbb{N}}$

$$\text{Ker}(\hat{S}) = \text{span}\{(1, 0, 0, \dots)\} \quad . \quad \text{Ker}(\hat{S}) = \{0\}$$

Thus $\text{Ind}(\hat{S}) = 1$

Index theorems connect index to a topological invariant

Structure: Toeplitz extension (no disorder)

S bilateral shift on $\ell^2(\mathbb{Z})$, then $C^*(S) \cong C(\mathbb{S}^1)$

\hat{S} unilateral shift on $\ell^2(\mathbb{N})$, only partial isometry with a defect:

$$\hat{S}^* \hat{S} = \mathbf{1} \quad \hat{S} \hat{S}^* = \mathbf{1} - |0\rangle\langle 0|$$

Then $C^*(\hat{S}) = \mathcal{T}$ Toeplitz algebra with exact sequence:

$$0 \rightarrow \mathcal{K} \xrightarrow{i} \mathcal{T} \xrightarrow{\pi} C(\mathbb{S}^1) \rightarrow 0$$

K -groups for any C^* -algebra \mathcal{A} (only rough definition):

$$K_0(\mathcal{A}) = \{[P] - [Q] : \text{projections in some } M_n(\mathcal{A})\}$$

$$K_1(\mathcal{A}) = \{[U] : \text{unitary in some } M_n(\mathcal{A})\}$$

Abelian group operation: Whitney sum

Example: $K_0(\mathbb{C}) = \mathbb{Z} = K_0(\mathcal{K})$ with invariant $\dim(P)$

Example: $K_1(C(\mathbb{S}^1)) = \mathbb{Z}$ with invariant given by winding number

6-term exact sequence for Toeplitz extension

C*-algebra short exact sequence \implies K-theory 6-term sequence

$$\begin{array}{ccccc} K_0(\mathcal{K}) = \mathbb{Z} & \xrightarrow{i_*} & K_0(T) = \mathbb{Z} & \xrightarrow{\pi_*} & K_0(\mathcal{C}(\mathbb{S}^1)) = \mathbb{Z} \\ \uparrow \text{Ind} & & & & \downarrow \text{Exp} \\ K_1(\mathcal{C}(\mathbb{S}^1)) = \mathbb{Z} & \xleftarrow{\pi_*} & K_1(T) = 0 & \xleftarrow{i_*} & K_1(\mathcal{K}) = 0 \end{array}$$

Here: $[A]_1 \in K_1(\mathcal{C}(\mathbb{S}^1))$ and $[\hat{P}\sigma_3]_0 = [\hat{P}_+]_0 - [\hat{P}_-]_0 \in K_0(\mathcal{K})$

$$\text{Ind}([A]_1) = [\hat{P}_+]_0 - [\hat{P}_-]_0 \quad (\text{bulk-boundary for } K\text{-theory})$$

$$\text{Ch}_0(\text{Ind}(A)) = \text{Ch}_1(A) \quad (\text{bulk-boundary for invariants})$$

Disordered case: analogous

4 K -theory crash course [RLL, WO]

K -theory developed to classify vector bundles over topological space X

Swan-Serre Theorem: $\{\text{vector bundles}\} \cong \{\text{projections in } M_n(C(X))\}$

Replace $C(X)$ by non-commutative C^* -algebra \mathcal{A}

Definition 4.1

$(\mathcal{A}, +, \cdot, \|\cdot\|)$ Banach algebra over \mathbb{C} if $\|AB\| \leq \|A\| \|B\|$, etc.

Then: \mathcal{A} is C^* -algebra $\iff \|A^*A\| = \|A\|^2$

Gelfand: commutative C^* algebras are $\mathcal{A} = C_0(X)$ with spectrum X

GNS: For any state on \mathcal{A} \exists Hilbert \mathcal{H} and irrep $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$

Example 1: $\mathcal{A} = \mathbb{C}$ or $\mathcal{A} = M_n(\mathbb{C})$

Example 2: Calkin's exact sequence over a Hilbert space \mathcal{H} :

$$0 \rightarrow \mathcal{K}(\mathcal{H}) \xrightarrow{i} \mathcal{B}(\mathcal{H}) \xrightarrow{\pi} \mathcal{Q}(\mathcal{H}) = \mathcal{B}(\mathcal{H})/\mathcal{K}(\mathcal{H}) \rightarrow 0$$

Definition of $K_0(\mathcal{A})$

Unitization $\mathcal{A}^+ = \mathcal{A} \oplus \mathbb{C}$ of C^* -algebra \mathcal{A} by

$$(\mathbf{A}, t)(\mathbf{B}, s) = (\mathbf{AB} + \mathbf{As} + \mathbf{Bt}, ts) \quad , \quad (\mathbf{A}, t)^* = (\mathbf{A}^*, \bar{t})$$

Natural C^* -norm $\|(\mathbf{A}, t)\| = \max\{\|\mathbf{A}\|, |t|\}$. Unit $\mathbf{1} = (0, 1) \in \mathcal{A}^+$

Exact sequence of C^* -algebras $0 \rightarrow \mathcal{A} \xrightarrow{i} \mathcal{A}^+ \xrightarrow{\rho} \mathbb{C} \rightarrow 0$

ρ has inverse $i'(t) = (0, t)$, then $s = i' \circ \rho : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ scalar part

$$\mathcal{V}_0(\mathcal{A}) = \left\{ V \in \bigcup_{n \geq 1} M_{2n}(\mathcal{A}^+) : V^* = V, V^2 = \mathbf{1}, s(V) \sim_0 E_{2n} \right\}$$

where $s(V) \sim_0 E_{2n}$ means homotopic to $E_{2n} = E_2^{\oplus n}$ with $E_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

Equivalence relation \sim_0 on $\mathcal{V}_0(\mathcal{A})$ by homotopy and $V \sim_0 \begin{pmatrix} V & 0 \\ 0 & E_2 \end{pmatrix}$

Then $K_0(\mathcal{A}) = \mathcal{V}_0(\mathcal{A}) / \sim_0$ abelian group via $[V]_0 + [V']_0 = \left[\begin{pmatrix} V & 0 \\ 0 & V' \end{pmatrix} \right]_0$

Definition of $K_0(\mathcal{A})$ is equivalent to standard one via $V = 2P - \mathbf{1}$:

$$K_0(\mathcal{A}) \cong \widehat{K}_0(\mathcal{A}) = \{[P] - [s(P)] : \text{projections in some } M_n(\mathcal{A}^+)\}$$

Satz 4.2 (Stability of K_0)

$$K_0(\mathcal{A}) = K_0(M_n(\mathcal{A})) = K_0(\mathcal{A} \otimes \mathcal{K})$$

Example 1: $K_0(\mathbb{C}) = K_0(\mathcal{K}) = \mathbb{Z}$, invariant $\dim(P) = \dim(\text{Ker}(V - \mathbf{1}))$

Example 2: $K_0(\mathcal{B}(\mathcal{H})) = 0$ for every separable \mathcal{H} by [RLL] 3.3.3

Example 3: $K_0(C(\mathbb{S}^1)) = \mathbb{Z}$ and $K_0(\mathcal{T}) = \mathbb{Z}$ for Toeplitz (also dim)

Satz 4.3 (0-cocycles paired with $K_0(\mathcal{A})$)

If \mathcal{T} tracial state on \mathcal{A} , then class map $\mathcal{T} : K_0(\mathcal{A}) \rightarrow \mathbb{R}$ defined by

$$\mathcal{T}[V]_0 = \mathcal{T}(P) = \frac{1}{2} \mathcal{T}(V - \mathbf{1})$$

Definition of $K_1(\mathcal{A})$

For definition of $K_1(\mathcal{A})$ set

$$\mathcal{V}_1(\mathcal{A}) = \left\{ U \in \bigcup_{n \geq 1} M_n(\mathcal{A}^+) : U^{-1} = U^* \right\}$$

Equivalence relation \sim_1 by homotopy and $U \sim_1 \begin{pmatrix} U & 0 \\ 0 & 1 \end{pmatrix}$

Then $K_1(\mathcal{A}) = \mathcal{V}_1(\mathcal{A}) / \sim_1$ with addition $[U]_1 + [U']_1 = [U \oplus U']_1$

If \mathcal{A} unital, one can work with $M_n(\mathcal{A})$ instead of $M_n(\mathcal{A}^+)$ in $\mathcal{V}_1(\mathcal{A})$

Example 1: $K_1(\mathbb{C}) = K_1(\mathcal{K}) = 0$

Example 2: $K_1(\mathcal{C}(\mathbb{S}^1)) = \mathbb{Z}$ with invariant "winding number"

Example 3: $K_1(\mathcal{A}^+) = K_1(\mathcal{A})$

Example 4: $K_1(\mathcal{B}(\mathcal{H})) = 0$ by Kuipers' theorem (holds for all W^* 's)

Example 5: For Calkin $K_1(\mathcal{Q}(\mathcal{H})) = \mathbb{Z}$ with invariant index of Fredholm

Suspension and Bott map

Definition 4.4

Suspension of a C^* -algebra \mathcal{A} is the C^* -algebra $S\mathcal{A} = C_0(\mathbb{R}) \otimes \mathcal{A}$

Alternatively upon rescaling: $S\mathcal{A} \cong C_0((0, 1), \mathcal{A})$

Satz 4.5 (Suspension)

One has isomorphism $\theta : K_1(\mathcal{A}) \rightarrow K_0(S\mathcal{A})$

Theorem 4.6 (Bott map)

One has isomorphism $\beta : K_0(\mathcal{A}) \cong \widehat{K}_0(\mathcal{A}) \rightarrow K_1(S\mathcal{A})$ given by

$$\beta([P]_0 - [s(P)]_0) = [t \in (0, 1) \mapsto (\mathbf{1} - P) + e^{2\pi it} P]_1$$

Note that r.h.s. indeed a unitary in $(S\mathcal{A})^+$

Korollar 4.7 (Bott periodicity)

$$K_0(SS\mathcal{A}) = K_0(\mathcal{A})$$

Construction of $\theta^{-1} : K_0(\mathcal{SA}) \rightarrow K_1(\mathcal{A})$ with adiabatic evolution:

$$0 \longrightarrow \mathcal{SA} \xrightarrow{i} C(\mathbb{S}^1, \mathcal{A}) \xrightarrow{\text{ev}} \mathcal{A} \longrightarrow 0$$

After rescaling is given a loop $t \in [0, 2\pi) \mapsto P_t = \frac{1}{2}(V_t + \mathbf{1}) \in M_N(\mathcal{A})$

With P_0 viewed as constant loop, $[P]_0 - [P_0]_0 \in K_0(\mathcal{SA})$

Indeed $\text{ev}([P]_0 - [P_0]_0) = 0$ so identified with element in $K_0(\mathcal{SA})$

Aim: find preimage under θ in $K_1(\mathcal{A})$

For $H_t = H_t^* \in M_N(\mathcal{A})$ satisfying $[H_t, P_t] = 0$ unitary solution $U_t \in \mathcal{A}^+$ of

$$i \partial_t U_t = (H_t + i[\partial_t P_t, P_t]) U_t, \quad U_0 = \mathbf{1}_N$$

Then $P_t = U_t P_0 U_t^*$ and $U_{2\pi} P_0 U_{2\pi}^* = P_0$

$$\theta^{-1}([P]_0 - [P_0]_0) = [P_0 U_{2\pi} P_0 + \mathbf{1}_N - P_0]_1$$

R.h.s. is unitary! Choice of H_t determines lift. Details in [PS] □

Natural push-forwards maps in K -theory

Associated to an exact sequence of C^* -algebras

$$0 \rightarrow \mathcal{K} \xrightarrow{i} \mathcal{A} \xrightarrow{\pi} \mathcal{Q} \rightarrow 0$$

there are natural push-forward maps:

$$i_* : K_j(\mathcal{K}) \rightarrow K_j(\mathcal{A}) \quad , \quad \pi_* : K_j(\mathcal{A}) \rightarrow K_j(\mathcal{Q})$$

given $i_*[V]_0 = [i(V)]_0$, $\pi_*[V]_0 = [\pi(V)]_0$, etc.

$\text{Ker}(\pi_*) = \text{Ran}(i_*)$, so short exact sequences of abelian groups:

$$K_0(\mathcal{K}) \xrightarrow{i_*} K_0(\mathcal{A}) \xrightarrow{\pi_*} K_0(\mathcal{Q})$$

and

$$K_1(\mathcal{Q}) \xleftarrow{\pi_*} K_1(\mathcal{A}) \xleftarrow{i_*} K_1(\mathcal{K})$$

Connecting maps close diagram to a cyclic 6-term diagram

Connecting maps from $K_j(\mathcal{Q})$ to $K_{j+1}(\mathcal{K})$

Definition 4.8 (Exponential map: $K_0(\mathcal{Q}) \rightarrow K_1(\mathcal{K})$)

Let $B = B^* \in M_n(\mathcal{A}^+)$ be contraction lift of unitary $V = V^* \in M_n(\mathcal{Q}^+)$

$$\begin{aligned}\text{Exp}[V]_0 &= [\exp(2\pi i(\frac{1}{2}(B + \mathbf{1})))]_1 \\ &= [-\cos(\pi B) - i \sin(\pi B)]_1 \\ &= [B\sqrt{\mathbf{1} - B^2} + i(\mathbf{1} - 2B^2)]_1\end{aligned}$$

Definition 4.9 (Index map: $K_1(\mathcal{Q}) \rightarrow K_0(\mathcal{K})$)

Let $B \in M_n(\mathcal{A}^+)$ be contraction lift of unitary $U \in M_n(\mathcal{Q}^+)$, namely $\pi^+(B) = U$ and $\|B\| \leq 1$. Then define

$$\text{Ind}[U]_1 = \left[\begin{pmatrix} 2BB^* - \mathbf{1} & 2B\sqrt{\mathbf{1} - B^*B} \\ 2B^*\sqrt{\mathbf{1} - BB^*} & \mathbf{1} - 2B^*B \end{pmatrix} \right]_0$$

Index map versus index of Fredholm operator

B unitary up to compact on $\mathcal{H} \iff \mathbf{1} - B^*B, \mathbf{1} - BB^* \in \mathcal{K}(\mathcal{H})$

$\implies B$ Fredholm operator and $U = \pi(B) \in \mathcal{Q}(\mathcal{H})$ unitary

Fedosov formula if $\mathbf{1} - B^*B$ and $\mathbf{1} - BB^*$ are traceclass:

$$\begin{aligned}\text{Ind}(B) &= \dim(\text{Ker}(B)) - \dim(\text{Ker}(B^*)) \\ &= \text{Tr}(\mathbf{1} - B^*B) - \text{Tr}(\mathbf{1} - BB^*) \\ &= \text{Tr} \begin{pmatrix} BB^* - \mathbf{1} & B(\mathbf{1} - B^*B)^{\frac{1}{2}} \\ (\mathbf{1} - B^*B)^{\frac{1}{2}} B^* & \mathbf{1} - B^*B \end{pmatrix} \\ &= \text{Tr}(\frac{1}{2}(V - \mathbf{1})) \\ &= \frac{1}{2} \text{Sig}(V) && \text{if } \mathbf{1} - B^*B, \mathbf{1} - BB^* \text{ projections} \\ &= \text{Tr}(\frac{1}{2}(\text{Ind}[U]_1 - \mathbf{1})) \\ &= \text{Tr}(\text{Ind}^\sim[U]_1)\end{aligned}$$

if $\text{Ind}^\sim[U]_1$ is the projection-valued version of index map

6-term exact sequence

Theorem 4.10

For every $0 \rightarrow \mathcal{K} \xrightarrow{i} \mathcal{A} \xrightarrow{\pi} \mathcal{Q} \rightarrow 0$, above definitions lead to

$$\begin{array}{ccccc} K_0(\mathcal{K}) & \xrightarrow{i_*} & K_0(\mathcal{A}) & \xrightarrow{\pi_*} & K_0(\mathcal{Q}) \\ \uparrow \text{Ind} & & & & \downarrow \text{Exp} \\ K_1(\mathcal{Q}) & \xleftarrow{\pi_*} & K_1(\mathcal{A}) & \xleftarrow{i_*} & K_1(\mathcal{K}) \end{array}$$

Proof in the books...

Example 4.11

Toeplitz extension $0 \rightarrow \mathcal{K}(\ell^2(\mathbb{N})) \xrightarrow{i} \mathcal{T} \xrightarrow{\pi} \mathcal{C}(\mathbb{S}^1) \rightarrow 0$

Bilateral shift $S \in \mathcal{C}(\mathbb{S}^1)$ gives class $[S]_1 \in K_1(\mathcal{C}(\mathbb{S}^1))$

Contraction lift is unilateral shift $\hat{S} \in \mathcal{T} \subset \mathcal{B}(\ell^2(\mathbb{N}))$ with $\hat{S}\hat{S}^* = \mathbf{1} - P_0$

From definition $\text{Ind}[S]_1 = [\text{diag}(\mathbf{1} - 2P_0, -\mathbf{1})]_0$

Exact sequence of the sphere

$$0 \rightarrow \mathbb{D}^{d+1} \hookrightarrow \overline{\mathbb{D}^{d+1}} \xrightarrow{\pi} \mathbb{S}^d \rightarrow 0$$

leads to an exact sequence of C^* -algebras

$$0 \rightarrow C_0(\mathbb{D}^{d+1}) \cong C_0(\mathbb{R}^{d+1}) \xrightarrow{i} C(\overline{\mathbb{D}^{d+1}}) \xrightarrow{\pi} C(\mathbb{S}^d) \rightarrow 0$$

All K -groups are well-known [WO]. For $d = 2n + 1$ odd

$$\begin{array}{ccccc}
 \mathbb{Z} & \xrightarrow{i_*} & \mathbb{Z} & \xrightarrow{\pi_*} & \mathbb{Z} \\
 \uparrow \text{Ind} & & & & \downarrow \text{Exp} \\
 \mathbb{Z} & \xleftarrow{\pi_*} & 0 & \xleftarrow{i_*} & 0
 \end{array}$$

while for $d = 2n$ even

$$\begin{array}{ccccc}
 0 & \xrightarrow{i_*} & \mathbb{Z} & \xrightarrow{\pi_*} & \mathbb{Z}^2 \\
 \uparrow \text{Ind} & & & & \downarrow \text{Exp} \\
 0 & \xleftarrow{\pi_*} & 0 & \xleftarrow{i_*} & \mathbb{Z}
 \end{array}$$

Aim: analyze one of the connecting maps, say Ind for d odd

Bott element

Let us write out $\text{Ind} : K_1(C(S^{2n-1})) = \mathbb{Z} \rightarrow K_0(C_0(\mathbb{D}^{2n})) = \mathbb{Z}$

For $n = 1$, generator is function $z : S^1 \rightarrow S^1$ with unit winding number

Lift is $z : \overline{\mathbb{D}^1} \rightarrow \overline{\mathbb{D}^1}$ which is *not* invertible, but a contraction

Bott element is "the" non-trivial projection on \mathbb{D}^2 :

$$\text{Ind}([z]_1) = \left[\begin{pmatrix} 2|z|^2 - 1 & 2z\sqrt{1-|z|^2} \\ 2\bar{z}\sqrt{1-|z|^2} & 1 - 2|z|^2 \end{pmatrix} \right]_0 \in K_0(C(\mathbb{D}^2))$$

For higher odd d , irrep $\gamma_1, \dots, \gamma_d$ of Clifford \mathbb{C}_d . Generator of $K_1(S^d)$

$$U = \sum_{j=1, \dots, d} x_j \gamma_j + i x_{d+1} \quad , \quad x = (x_1, \dots, x_{d+1}) \in S^d$$

Lift $B \in C(\overline{\mathbb{D}^{d+1}})$ same formula. Then with $r = \|x\|$

$$\text{Ind}[U]_1 = \left[\begin{pmatrix} 2r^2 - 1 & 2(1 - r^2)^{\frac{1}{2}} B \\ 2B^*(1 - r^2)^{\frac{1}{2}} & -(2r^2 - 1) \end{pmatrix} \right]_0$$

Fuzzy spheres and their index map

Definition 4.12

A unital C^* -algebra. A fuzzy d -sphere of width $\delta < 1$ is a collection of self-adjoints $X_1, \dots, X_{d+1} \in \mathcal{A}$ with spectrum in $[-1, 1]$ such that

$$\|\mathbf{1} - \sum_{j=1, \dots, d+1} (X_j)^2\| < \delta \quad , \quad \|[X_j, X_i]\| < \delta$$

Proposition 4.13

If d is odd, a fuzzy d -sphere in \mathcal{Q} specifies an element $[A]_1 \in K_1(\mathcal{Q})$ via

$$A = \sum_{j=1, \dots, d} X_j \gamma_j + i X_{d+1}$$

Theorem 4.14

For $0 \rightarrow \mathcal{K} \xrightarrow{i} \mathcal{A} \xrightarrow{\pi} \mathcal{Q} \rightarrow 0$ and with B is lift of A and $B^*B = R^2$

$$\text{Ind}[A]_1 = \left[\begin{pmatrix} 2R^2 - 1 & 2(1 - R^2)^{\frac{1}{2}} B \\ 2B^*(1 - R^2)^{\frac{1}{2}} & -(2R^2 - 1) \end{pmatrix} \right]_0$$

... and there is Real KR -theory (Karoubi,...)

Requires the data of a real structure in form of an anti-linear involution

$$\tau : \mathcal{A} \rightarrow \mathcal{A} \quad , \quad \tau(A + \lambda B) = \tau(A) + \bar{\lambda}\tau(B) \quad , \quad \tau^2 = \mathbf{1}$$

Then

$$\mathcal{V}_0^\tau(\mathcal{A}) = \left\{ V \in \cup_{n \geq 1} M_{2n}(\mathcal{A}^+) : V^* = V = \tau(V), V^2 = \mathbf{1}, s(V) \sim_0 E_{2n} \right\}$$

Now homotopy \sim_0 within $\mathcal{V}_0^\tau(\mathcal{A})$

Then $KR_0(\mathcal{A}) = \mathcal{V}_0^\tau(\mathcal{A}) / \sim_0$ abelian group via $[V]_0 + [V']_0 = \left[\begin{pmatrix} V & 0 \\ 0 & V' \end{pmatrix} \right]_0$

Going on from here there 7 further groups $KR_1(\mathcal{A}), \dots, KR_7(\mathcal{A})$

One has $KR_j(\mathcal{A}) = KR_0(S^j \mathcal{A})$ for Real suspension

Bott periodicity is $KR_0(\mathcal{A}) = KR_0(S^8 \mathcal{A})$

Hence KR -theory is 8 periodic

For exact sequence of Real algebras there is 64-term diagram ($= 8 \cdot 3$)

Here no further details because possibly not so important for physics

5 Observable algebra for tight-binding models

One-particle Hilbert space $\ell^2(\mathbb{Z}^d) \otimes \mathbb{C}^L$

Fiber $\mathbb{C}^L = \mathbb{C}^{2s+1} \otimes \mathbb{C}^r$ with spin s and r internal degrees

e.g. $\mathbb{C}^r = \mathbb{C}_{\text{ph}}^2 \otimes \mathbb{C}_{\text{sl}}^2$ particle-hole space and sublattice space

Typical Hamiltonian

$$H_\omega = \Delta^B + W_\omega = \sum_{i=1}^d (t_i^* S_i^B + t_i (S_i^B)^*) + W_\omega$$

Magnetic translations $S_j^B S_i^B = e^{iB_{i,j}} S_i^B S_j^B$ in Landau gauge:

$$S_1^B = S_1 \quad S_2^B = e^{iB_{1,2}X_1} S_2 \quad S_3^B = e^{iB_{1,3}X_1 + iB_{2,3}X_2} S_3$$

t_i matrices $L \times L$, e.g. spin orbit coupling, (anti)particle creation

matrix potential $W_\omega = W_\omega^* = \sum_{n \in \mathbb{Z}^d} |n\rangle \omega_n \langle n|$ with i.i.d. matrices ω_n

Configurations $\omega = (\omega_n)_{n \in \mathbb{Z}^d} \in \Omega$ compact probability space (Ω, \mathbb{P})

\mathbb{P} invariant and ergodic w.r.t. $T : \mathbb{Z}^d \times \Omega \rightarrow \Omega$

Covariant operators (generalizes periodicity)

Covariance w.r.t. to dual magnetic translations $V_a = S_j^B V_a (S_j^B)^*$

$$V_a H_\omega V_a^* = H_{T_a \omega} \quad , \quad a \in \mathbb{Z}^d$$

$\|A\| = \sup_{\omega \in \Omega} \|A_\omega\|$ is C*-norm on

$$\begin{aligned} \mathcal{A}_d &= C^* \{ A = (A_\omega)_{\omega \in \Omega} \text{ finite range covariant operators} \} \\ &\cong \text{twisted crossed product } C(\Omega) \rtimes_B \mathbb{Z}^d \end{aligned}$$

Fact: Suppose Ω contractible

\implies rotation algebra $C^*(S_1^B, \dots, S_d^B)$ is deformation retract of \mathcal{A}_d

In particular: K -groups of $C^*(S_1^B, \dots, S_d^B)$ and \mathcal{A}_d coincide

Theorem 5.1 (Pimsner-Voiculescu 1980)

$$K_0(\mathcal{A}_d) = \mathbb{Z}^{2^{d-1}} \text{ and } K_1(\mathcal{A}_d) = \mathbb{Z}^{2^{d-1}}$$

Generators of $K_j(\mathcal{A}_d)$

Pimsner-Voiculescu also showed that there are short exact sequences:

$$0 \rightarrow K_0(\mathcal{A}_{d-1}) \xrightarrow{i_*} K_0(\mathcal{A}_d) \xrightarrow{\text{Exp}} K_1(\mathcal{A}_{d-1}) \rightarrow 0$$

$$0 \rightarrow K_1(\mathcal{A}_{d-1}) \xrightarrow{i_*} K_1(\mathcal{A}_d) \xrightarrow{\text{Ind}} K_0(\mathcal{A}_{d-1}) \rightarrow 0$$

Both lines read $K_j(\mathcal{A}_d) = K_0(\mathcal{A}_{d-1}) \oplus K_1(\mathcal{A}_{d-1}) = \mathbb{Z}^{2^{d-2}} \oplus \mathbb{Z}^{2^{d-2}}$

Iterative construction of generators using inverse of Ind and Exp

Explicit generators $[G_I]$ of K -groups labelled by subsets $I \subset \{1, \dots, d\}$

Top generator $I = \{1, \dots, d\}$ identified with Bott in $K_j(C(\mathbb{S}^d))$

Example $G_{\{1,2\}}$ Powers-Rieffel projection and $C^*(S_1^B, S_2^B)$

In general, any projection $P \in M_n(\mathcal{A}_d)$ can be decomposed as

$$[P]_0 = \sum_{I \subset \{1, \dots, d\}} n_I [G_I]_0 \quad n_I \in \mathbb{Z}, |I| \text{ even}$$

Questions: calculate $n_I = c_I \text{Ch}_I(P)$, physical significance?

K -group elements of physical interest

Fermi level $\mu \in \mathbb{R}$ in spectral gap of H_ω

$$P_\omega = \chi(H_\omega \leq \mu) \quad \text{covariant Fermi projection}$$

Hence: $P = (P_\omega)_{\omega \in \Omega} \in \mathcal{A}_d$ fixes element in $[P]_0 \in K_0(\mathcal{A}_d)$

If chiral symmetry present: Fermi unitary $U = A|A|^{-1}$ from

$$H_\omega = -J_{\text{ch}}^* H_\omega J_{\text{ch}} = \begin{pmatrix} 0 & A_\omega \\ A_\omega^* & 0 \end{pmatrix}, \quad J_{\text{ch}} = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}$$

If $\mu = 0$ in gap, $A = (A_\omega)_{\omega \in \Omega} \in \mathcal{A}_d$ invertible and $[U]_1 = [A]_1 \in K_1(\mathcal{A}_d)$

Remark Sufficient to have an approximate chiral symmetry

$$H_\omega = \begin{pmatrix} B_\omega & A_\omega \\ A_\omega^* & C_\omega \end{pmatrix} \quad \text{with invertible } A_\omega$$

Strong and weak invariants

Fermi level $\mu \implies$ Fermi projection P or Fermi unitary A

Decompositions

$$[P]_0 = \sum_{I \subset \{1, \dots, d\}} n_I [G_I]_0 \quad , \quad [A]_1 = \sum_{I \subset \{1, \dots, d\}} n_I [G_I]_1$$

Invariants n_I , top invariant $n_{\{1, \dots, d\}} \in \mathbb{Z}$ called *strong*, others weak

A systems with $n_{\{1, \dots, d\}} \neq 0$ is called a strong topological insulator

If $n_{\{1, \dots, d\}} = 0$, but some other $n_I \neq 0$, weak topological insulator

For Class A (no symmetry) and Class AIII (chiral symmetry):

	dimension d	1	2	3	4	5	6	7	8
A	strong invariant	0	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}
AIII	strong invariant	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0	\mathbb{Z}	0

\mathbb{Z} -entries are parts of the K -groups. Calculation of number later

Non-commutative analysis tools [BES, PS]

Definition 5.2 (Non-commutative integration and derivatives)

Tracial state \mathcal{T} on \mathcal{A}_d given by

$$\mathcal{T}(A) = \mathbf{E}_{\mathbb{P}} \operatorname{Tr}_L \langle 0 | A_\omega | 0 \rangle$$

Derivations $\nabla = (\nabla_1, \dots, \nabla_d)$ densely defined by

$$\nabla_j A_\omega = i[X_j, A_\omega]$$

Then define $C^k(\mathcal{A})$, $C^\infty(\mathcal{A})$, etc.

Usual rules: $\mathcal{T}(AB) = \mathcal{T}(BA)$, $\nabla(AB) = \nabla(A)B + A\nabla(B)$, etc.

Also: $\mathcal{T}(\nabla(A)) = 0$, so partial integration $\mathcal{T}(\nabla(A)B) = -\mathcal{T}(A\nabla(B))$

Proposition 5.3 (Birkhoff theorem for translation group)

\mathcal{T} is \mathbb{P} -almost surely the trace per unit volume

$$\mathcal{T}(A) = \lim_{\Lambda \rightarrow \mathbb{Z}^d} \frac{1}{|\Lambda|} \sum_{n \in \Lambda} \operatorname{Tr}_L \langle n | A_\omega | n \rangle$$

Periodic systems

For simplicity 1-periodic in all directions and no magnetic field

Then $\mathcal{A}_d = C(\mathbb{T}^d) \otimes \mathbb{C}^{L \times L}$ commutative up to matrix degree

non-commutative	A	$\nabla_j(A)$	\mathcal{T}
commutative	$k \mapsto A(k)$	$\partial_{k_j} A$	$\int_{\mathbb{T}^d} dk \text{Tr}$

With dictionary: rewrite many formulas from solid state literature

Example: Kubo formula for conductivity at relaxation time τ

$$\int dk \sum_{n,m} \left(\partial_{k_j} f_{\beta,\mu}(E_n(k)) (E_n(k) - E_m(k) + \frac{1}{\tau})^{-1} \partial_{k_j} E_m(k) \right) \\ = \mathcal{T} \left(\nabla_i(f_{\beta,\mu}(H)) (\mathcal{L}_H + \frac{1}{\tau})^{-1} (\nabla_j(H)) \right)$$

where $\mathcal{L}_H = i[H, \cdot]$ Liouville operator

6 Topological invariants in solid state systems

For invertible $A \in \mathcal{A}_d$ and odd $|I|$, with $\rho : \{1, \dots, |I|\} \rightarrow I$:

$$\text{Ch}_I(A) = \frac{i(i\pi)^{\frac{|I|-1}{2}}}{|I|!!} \sum_{\rho \in \mathcal{S}_I} (-1)^\rho \mathcal{T} \left(\prod_{j=1}^{|I|} A^{-1} \nabla_{\rho_j} A \right) \in \mathbb{R}$$

where $\mathcal{T}(A) = \mathbf{E}_{\mathbb{P}} \text{Tr}_L \langle 0 | A_\omega | 0 \rangle$ and $\nabla_j A_\omega = i[X_j, A_\omega]$

For even $|I|$ and projection $P \in \mathcal{A}_d$:

$$\text{Ch}_I(P) = \frac{(2i\pi)^{\frac{|I|}{2}}}{\frac{|I|}{2}!} \sum_{\rho \in \mathcal{S}_I} (-1)^\rho \mathcal{T} \left(P \prod_{j=1}^{|I|} \nabla_{\rho_j} P \right) \in \mathbb{R}$$

Theorem 6.1 (Connes 1985, [Con])

$\text{Ch}_I(A)$ and $\text{Ch}_I(P)$ homotopy invariants; pairings with $K(\mathcal{A}_d)$

Link to Volovik-Essin-Gurarie invariants

Express the invariants in terms of Green function/resolvent

Consider path $z : [0, 1] \rightarrow \mathbb{C} \setminus \sigma(H)$ encircling $(-\infty, \mu] \cap \sigma(H)$

Set

$$G(t) = (H - z(t))^{-1}$$

Theorem 6.2 ([PS])

For $|l|$ even and with $\nabla_0 = \partial_t$,

$$\text{Ch}_l(P_\mu) = \frac{(i\pi)^{\frac{|l|}{2}}}{i(|l| - 1)!!} \sum_{\rho \in \mathcal{S}_{l \cup \{0\}}} (-1)^\rho \int_0^1 dt \mathcal{T} \left(\prod_{j=0}^{|l|} G(t)^{-1} \nabla_{\rho_j} G(t) \right)$$

Isomorphism via Bott map $\beta : K_0(\mathcal{A}_d) \rightarrow K_1(\mathcal{S}\mathcal{A}_d)$ leads to

$$\beta[P_\mu]_0 = [t \in [0, 1] \mapsto G(t)]_1$$

Combine with suspension result on cyclic cohomology side

Similar results for odd pairings

Generalized Streda formulæ

In QHE: integrated density of states grows linearly in magnetic field

integrated density of states: $\mathbf{E} \langle 0|P|0\rangle = \text{Ch}_\emptyset(P)$

$$\partial_{B_{1,2}} \text{Ch}_\emptyset(P) = \frac{1}{2\pi} \text{Ch}_{\{1,2\}}(P)$$

Theorem 6.3 ([PS])

$$\partial_{B_{i,j}} \text{Ch}_I(P) = \frac{1}{2\pi} \text{Ch}_{I \cup \{i,j\}}(P) \quad |I| \text{ even, } i, j \notin I$$

$$\partial_{B_{i,j}} \text{Ch}_I(A) = \frac{1}{2\pi} \text{Ch}_{I \cup \{i,j\}}(A) \quad |I| \text{ odd, } i, j \notin I$$

Application: magneto-electric effects in $d = 3$

Time is 4th direction needed for calculation of polarization

Non-linear response is derivative w.r.t. B given by $\text{Ch}_{\{1,2,3,4\}}(P)$

Index theorem for strong invariants and odd d

$\gamma_1, \dots, \gamma_d$ irrep of Clifford C_d on $\mathbb{C}^{2^{(d-1)/2}}$

$$D = \sum_{j=1}^d X_j \otimes \mathbf{1} \otimes \gamma_j \quad \text{Dirac operator on } \ell^2(\mathbb{Z}^d) \otimes \mathbb{C}^L \otimes \mathbb{C}^{2^{(d-1)/2}}$$

Dirac phase $F = \frac{D}{|D|}$ provides odd Fredholm module on \mathcal{A}_d :

$$F^2 = \mathbf{1} \quad [F, A_\omega] \text{ compact and in } \mathcal{L}^{d+\epsilon} \text{ für } A = (A_\omega)_{\omega \in \Omega} \in \mathcal{A}_d$$

Theorem 6.4 (Local index = generalizes Noether-Gohberg-Krein)

Let $\Pi = \frac{1}{2}(F + \mathbf{1})$ be Hardy Projektion for F . For invertible A_ω

$$\text{Ch}_{\{1, \dots, d\}}(A) = \text{Ind}(\Pi A_\omega \Pi)$$

The index is \mathbb{P} -almost surely constant.

Local index theorem for even dimension d

As above $\gamma_1, \dots, \gamma_d$ Clifford, grading $\Gamma = -i^{-d/2} \gamma_1 \cdots \gamma_d$

Dirac $D = -\Gamma D \Gamma = |D| \begin{pmatrix} 0 & F \\ F^* & 0 \end{pmatrix}$ even Fredholm module

Theorem 6.5 (Connes $d = 2$, Prodan, Leung, Bellissard 2013)

Almost sure index $\text{Ind}(P_\omega F P_\omega)$ equal to $\text{Ch}_{\{1, \dots, d\}}(P)$

Special case $d = 2$: $F = \frac{X_1 + iX_2}{|X_1 + iX_2|}$ and

$$\text{Ind}(P_\omega F P_\omega) = 2\pi i \mathcal{T}(P[[X_1, P], [X_2, P]])$$

Proofs: geometric identity of high-dimensional simplex

Advantages: phase label also for dynamical localized regime
implementation of discrete symmetries (CPT)

Bott operator (Loring and Hastings 2008, [Lor, LSB])

For tuning parameter $\kappa > 0$ and invertible local A :

$$B_\kappa = \begin{pmatrix} \kappa D & A \\ A^* & -\kappa D \end{pmatrix} = \kappa D \otimes \sigma_3 + H$$

where $H = \begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix}$ chiral Hamiltonian. Clearly B_κ selfadjoint

D unbounded with discrete spectrum, A viewed as perturbation

A may lead to spectral asymmetry of B_κ , but not for $A = \mathbf{1}$

Measured by signature, already on finite volume approximation!

A_ρ restriction of A (Dirichlet b.c.) to $\mathbb{D}_\rho = \{x \in \mathbb{Z}^d : |x| \leq \rho\}$

$$B_{\kappa,\rho} = \begin{pmatrix} \kappa D_\rho & A_\rho \\ A_\rho^* & -\kappa D_\rho \end{pmatrix}$$

Finite volume calculation of topological invariants

Theorem 6.6 ([LSB])

Let $g = \|A^{-1}\|^{-1}$ be the invertibility gap. Provided that

$$\|[D, A]\| \leq \frac{g^3}{18 \|A\| \kappa} \quad (*)$$

and

$$\frac{2g}{\kappa} \leq \rho \quad (**)$$

the matrix $B_{\kappa, \rho}$ is invertible and strong invariant is

$$\frac{1}{2} \text{Sig}(B_{\kappa, \rho}) = \text{Ind}(\Pi A \Pi + (\mathbf{1} - \Pi))$$

How to use: form (*) infer κ , then ρ from (**)

If A unitary, $g = \|A\| = 1$ and $\kappa = (18\|[D, A]\|)^{-1}$ and $\rho = 2/\kappa$

Hence **small** matrix of size ≤ 100 sufficient! Great for numerics!

Why it can work:

Proposition 6.7

If (*) and (**) hold (which includes the case $\rho = \infty$),

$$B_{\kappa,\rho}^2 \geq \frac{g^2}{2}$$

Proof:

$$B_{\kappa,\rho}^2 = \begin{pmatrix} A_\rho^* A_\rho & 0 \\ 0 & A_\rho A_\rho^* \end{pmatrix} + \kappa^2 \begin{pmatrix} D_\rho^2 & 0 \\ 0 & D_\rho^2 \end{pmatrix} + \kappa \begin{pmatrix} 0 & [D_\rho, A_\rho] \\ [D_\rho, A_\rho]^* & 0 \end{pmatrix}$$

Last term is a perturbation controlled by (*)

First two terms positive (indeed: close to origin and away from it)

Now $A^*A \geq g^2$, but $(A^*A)_\rho \neq A_\rho^*A_\rho$

This issue can be dealt with by tapering argument:

Proposition 6.8 (Bratelli-Robinson)

For $f : \mathbb{R} \rightarrow \mathbb{R}$ with Fourier transform defined without $\sqrt{2\pi}$,

$$\|[f(D), A]\| \leq \|\hat{f}'\|_1 \|[D, A]\|$$

Lemma 6.9 (Tapering function)

\exists even function $f : \mathbb{R} \rightarrow [0, 1]$ with $f(x) = 0$ for $|x| \geq \rho$
and $f(x) = 1$ for $|x| \leq \frac{\rho}{2}$ such that $\|\hat{f}'\|_1 = \frac{8}{\rho}$

With this, $f = f(D) = f(|D|)$ and $\mathbf{1}_\rho = \chi(|D| \leq \rho)$:

$$\begin{aligned} A_\rho^* A_\rho &= \mathbf{1}_\rho A^* \mathbf{1}_\rho A \mathbf{1}_\rho \geq \mathbf{1}_\rho A^* f^2 A \mathbf{1}_\rho \\ &= \mathbf{1}_\rho f A^* A f \mathbf{1}_\rho + \mathbf{1}_\rho ([A^*, f] f A + f A^* [f, A]) \mathbf{1}_\rho \\ &\geq g^2 f^2 + \mathbf{1}_\rho ([A^*, f] f A + f A^* [f, A]) \mathbf{1}_\rho \end{aligned}$$

So indeed $A_\rho^* A_\rho$ positive close to origin

Then one can conclude... but TEDIOUS □

η -invariant (Atiyah-Patodi-Singer 1977)

Definition 6.10

$B = B^*$ invertible operator on \mathcal{H} with compact resolvent. Then

$$\eta(B) = \text{Tr}(B|B|^{-s-1})|_{s=0} = \frac{1}{\Gamma(\frac{s+1}{2})} \int_0^\infty dt t^{\frac{s-1}{2}} \text{Tr}(B e^{-tB^2})|_{s=0}$$

provided it exists!

If $\dim(\mathcal{H}) < \infty$, then $\eta(B) = \text{Sig}(B)$

Usually existence of η -invariant for ψ -Diffs difficult issue

Proposition 6.11

If (*) holds, B_κ has well-defined η -invariant

Proof. Integral for large t controlled by gap (Proposition above)

For small t appeal to Dyson series (iteration of DuHamel):

$$e^{-tB_\kappa^2} = e^{-t\Delta} + t \int_0^1 dr e^{-(1-r)t\Delta} R e^{-rtB_\kappa^2}$$

where $B_\kappa^2 = \Delta + R$ with

$$\Delta = \kappa^2 \begin{pmatrix} D^2 & 0 \\ 0 & D^2 \end{pmatrix}, \quad R = \begin{pmatrix} AA^* & \kappa[D, A] \\ \kappa[D, A]^* & A^*A \end{pmatrix}$$

Now replacing $B_\kappa = \kappa D \otimes \sigma_3 + H$

$$\text{Tr}(B_\kappa e^{-t\Delta}) = \kappa \text{Tr} \left(\begin{pmatrix} D & 0 \\ 0 & -D \end{pmatrix} e^{-t\Delta} \right) + \text{Tr} \left(\begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix} e^{-t\Delta} \right) = 0$$

Second term has supplementary factor t



Theorem 6.12 (follows from Getzler 1993, Carey-Phillips 2004)

Suppose (*) so that B_κ has well-defined η -invariant

For path $\lambda \in [0, 1] \mapsto B_\kappa(\lambda) = \kappa D \otimes \sigma_3 + \lambda H$ of selfadjoints

$$2 \text{SF}(\lambda \in [0, 1] \mapsto B_\kappa(\lambda)) = \eta(B_\kappa(1)) - \eta(B_\kappa(0)) = \eta(B_\kappa)$$

Consequence: As spectral flow homotopy invariant, so is $\eta(B_\kappa)$

Using this, **first proof** of Theorem 6.6 for dimension $d = 1$:

By homotopy invariance sufficient: $A = S^n$ for $n \in \mathbb{Z}$ and S shift

Then calculate spectrum of $B_\kappa(\lambda)$ explicitly using $XS = (X + 1)S$:

$$\sigma(B_\kappa(\lambda)) = \left\{ \frac{\kappa}{2} \left(n \pm \left((n - 2k)^2 + \frac{4\lambda^2}{\kappa^2} \right)^{\frac{1}{2}} \right) : k \in \mathbb{Z} \right\}$$

Now carefully follow eigenvalues to calculate spectral flow □

Localizing index map for index pairings

Suppose now $U = \pi(\Pi A \Pi + (\mathbf{1} - \Pi)) \in \mathcal{Q}$ as in Theorem 6.6

but first A unitary. Then contraction lift $B = \Pi A \Pi + (\mathbf{1} - \Pi)$

Modify Π and $\mathbf{1} - \Pi$ to $\rho = \rho(D)$ smooth and $n = n(D)$ where

$$\rho(x) = \begin{cases} 0, & x \leq -\rho \\ \rho(x), & |x| \leq \rho \\ 1, & x \geq \rho \end{cases}, \quad n(x) = \begin{cases} 1, & x \leq -\rho \\ 0, & x \geq \rho \end{cases}$$

Now $\rho - \Pi$, $n - (\mathbf{1} - \Pi)$ compact, $n\rho = \rho n = 0$ and $n + \rho|_{\mathbb{D}_\rho^c} = \mathbf{1}_{\mathbb{D}_\rho^c}$

With notation $A_\rho = \rho A \rho$ acting only on $\ell^2(\mathbb{D}_\rho) \otimes \mathbb{C}^N$:

$$\begin{aligned} \text{Ind}[U] &= \text{Ind}[\rho A \rho + n] = \text{Ind}[A_\rho + n] \\ &= \left[\begin{pmatrix} 2A_\rho A_\rho^* - \mathbf{1} & 2A_\rho(\mathbf{1} - A_\rho^* A_\rho)^{\frac{1}{2}} \\ 2(\mathbf{1} - A_\rho^* A_\rho)^{\frac{1}{2}} A_\rho^* & \mathbf{1} - 2A_\rho^* A_\rho \end{pmatrix} \oplus \begin{pmatrix} \mathbf{1}_{\mathbb{D}_\rho^c} & 0 \\ 0 & -\mathbf{1}_{\mathbb{D}_\rho^c} \end{pmatrix} \right] \end{aligned}$$

Summand on \mathbb{D}_ρ^c trivial (as equal to E_2). Thus:

$$\text{Ind}[U] = \left[\begin{pmatrix} 2A_\rho A_\rho^* - \mathbf{1} & 2A_\rho(\mathbf{1} - A_\rho^* A_\rho)^{\frac{1}{2}} \\ 2(\mathbf{1} - A_\rho^* A_\rho)^{\frac{1}{2}} A_\rho^* & \mathbf{1} - 2A_\rho^* A_\rho \end{pmatrix} \right]$$

Numerical index is signature of this finite-dimensional matrix!

Modify to self-adjoint matrix without spoiling invertibility

$$\begin{aligned} \|A_\rho A_\rho^* - p^4\| &= \|p A p^2 A^* p - p^3 A A^* p\| \leq \|[p^2, A]\| \\ &\leq \frac{C}{\rho} \|[D, A]\| < \frac{1}{4} \end{aligned}$$

by the smoothness of p and for ρ sufficiently large. Similarly

$$\|A_\rho(\mathbf{1} - A_\rho^* A_\rho)^{\frac{1}{2}} - (\mathbf{1} - p^4)^{\frac{1}{4}} p A p (\mathbf{1} - p^4)^{\frac{1}{4}}\| \leq \frac{C}{\rho} \|[D, A]\| < \frac{1}{4}$$

Thus just replace matrix entries without changing signature!

Proposition 6.13

If (*) and (**) hold,

$$\begin{aligned} & \text{Ind}(\Pi A \Pi + (\mathbf{1} - \Pi)) \\ &= \text{Sig} \begin{pmatrix} 2p^4 - \mathbf{1} & 2(\mathbf{1} - p^4)^{\frac{1}{4}} p A p (\mathbf{1} - p^4)^{\frac{1}{4}} \\ 2(\mathbf{1} - p^4)^{\frac{1}{4}} p A^* p (\mathbf{1} - p^4)^{\frac{1}{4}} & \mathbf{1} - 2p^4 \end{pmatrix} \end{aligned}$$

Last tasks:

1) replace $2p^4 - \mathbf{1}$ by κD_ρ

2) replace $\sqrt{2}(\mathbf{1} - p^4)^{\frac{1}{4}} p$ by $\mathbf{1}_\rho$ indicator on \mathbb{D}_ρ . Then $\mathbf{1}_\rho A \mathbf{1}_\rho = A_\rho$

Both follows again by a tapering argument

UUuuuffff

7 Invariants as response coefficients

- Hall conductance via Kubo formula: $\text{Ch}_{\{i,j\}}$ with $i \neq j$
- polarization for periodically driven systems: $\text{Ch}_{\{0,j\}}$ with 0 time
- orbital magnetization at zero temperature
- magneto-electric effect: $\text{Ch}_{\{0,1,2,3\}}$ with 0 time
- chiral polarization: $\text{Ch}_{\{j\}}$

Current operator $J = (J_1, \dots, J_d)$ in d dimension:

$$J = \dot{X} = i[H, X] = \nabla H$$

Current density at equilibrium expressed by Fermi-Dirac state:

$$j_{\beta,\mu} = \mathcal{T}(f_{\beta,\mu}(H) J) \quad , \quad f_{\beta,\mu}(H) = (\mathbf{1} + e^{\beta(H-\mu)})^{-1}$$

Proposition 7.1 ([BES])

If $H = H^* \in C^1(\mathcal{A})$ and $f \in C_0(\mathbb{R})$, then $\mathcal{T}(f(H)\nabla H) = 0$

Proof: Leibniz implies $0 = \mathcal{T}(\nabla H^n) = n\mathcal{T}(H^{n-1}\nabla H)$ for all $n \geq 1$ \square

Hence no current at equilibrium! Add external electric field $\mathcal{E} \in \mathbb{R}^d$

$$H_{\mathcal{E}} = H + \mathcal{E} \cdot X$$

Then $H_{\mathcal{E}}$ neither bounded nor homogeneous and thus not in \mathcal{A}

Nevertheless associated time evolution remains in the algebra \mathcal{A}

In the Schrödinger picture it is governed by the Liouville equation:

$$\partial_t \rho = -i[H_{\mathcal{E}}, \rho] = -i[H + \mathcal{E} \cdot X, \rho] = -\mathcal{L}_H(\rho) + \mathcal{E} \cdot \nabla(\rho)$$

Now Dyson series with Liouville \mathcal{L}_H as perturbation is iteration of

$$e^{t\mathcal{L}_{H_{\mathcal{E}}}} = e^{t\mathcal{E} \cdot \nabla} + \int_0^t ds e^{(t-s)\mathcal{E} \cdot \nabla} \mathcal{L}_H e^{s\mathcal{L}_{H_{\mathcal{E}}}}$$

This shows:

Proposition 7.2

$\pm \mathcal{L}_H + \mathcal{E} \cdot \nabla$ are generators of automorphism groups in \mathcal{A}

Next time-averaged current under the dynamics with \mathcal{E} :

$$j_{\beta,\mu,\mathcal{E}} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \mathcal{T}(f_{\beta,\mu}(H) e^{t\mathcal{L}_{H\mathcal{E}}}(J))$$

As trace \mathcal{T} invariant under both ∇ and \mathcal{L}_H ,

$$j_{\beta,\mu,\mathcal{E}} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \mathcal{T}(J e^{-t\mathcal{L}_{H\mathcal{E}}}(f_{\beta,\mu}(H)))$$

(Schrödinger picture \iff Heisenberg picture). Now

Proposition 7.3 (Bloch Oscillations)

Time-averaged current $j_{\beta,\mu,\mathcal{E}}$ along direction of \mathcal{E} vanishes

Proof. $\mathcal{E} \cdot J(t) = e^{t\mathcal{L}_{H\mathcal{E}}}(\mathcal{E} \cdot \nabla(H)) = e^{t\mathcal{L}_{H\mathcal{E}}}(\mathcal{L}_{H\mathcal{E}}(H)) = \frac{dH(t)}{dt}$

Taking the time average gives us

$$\frac{1}{T} \int_0^T dt \mathcal{E} \cdot J(t) = \frac{H(T) - H}{T}$$

Since H bounded and $\|H(t)\| = \|H\|$, r.h.s. vanishes as $T \rightarrow \infty$ □

Modify dynamics by bounded linear collision term (like Boltzmann eq.):

$$\partial_t \rho + \mathcal{L}_H(\rho) - \mathcal{E} \cdot \nabla(\rho) = -\Gamma(\rho)$$

Main property is invariance of equilibrium: $\Gamma(f_{\beta,\mu}(H)) = 0$

Again Dyson series shows existence of dynamics:

$$\rho(t) = e^{-t(\mathcal{L}_H - \mathcal{E} \cdot \nabla + \Gamma)}(\rho(0))$$

Initial state chosen to be $\rho(0) = f_{\beta,\mu}(H)$

Exponential time-averaged current density shows:

$$\begin{aligned} j_{\beta,\mu,\mathcal{E}} &= \lim_{\delta \rightarrow 0} \delta \int_0^\infty dt e^{-\delta t} \mathcal{T}(J\rho(t)) \\ &= \lim_{\delta \rightarrow 0} \delta \mathcal{T} \left(J \frac{1}{\delta + \Gamma + \mathcal{L}_H - \mathcal{E} \cdot \nabla} (f_{\beta,\mu}(H)) \right) \end{aligned}$$

By Proposition 7.1 and $(\mathcal{L}_H + \Gamma)(f_{\beta,\mu}(H)) = 0$ no current at equilibrium:

$$0 = \delta \mathcal{T} \left(J \frac{1}{\delta} f_{\beta,\mu}(H) \right) = \delta \mathcal{T} \left(J \frac{1}{\delta + \mathcal{L}_H + \Gamma} (f_{\beta,\mu}(H)) \right)$$

Subtract this from $j_{\beta,\mu,\mathcal{E}}$ and use resolvent identity

$$j_{\beta,\mu,\mathcal{E}} = \lim_{\delta \rightarrow 0} \mathcal{T} \left(\mathcal{J} \frac{1}{\delta + \Gamma + \mathcal{L}_H - \mathcal{E} \cdot \nabla} \mathcal{E} \cdot \nabla \frac{\delta}{\delta + \Gamma + \mathcal{L}_H} (f_{\beta,\mu}(H)) \right)$$

Now, again $(\mathcal{L}_H + \Gamma)(f_{\beta,\mu}(H)) = 0$,

$$j_{\beta,\mu,\mathcal{E}} = \lim_{\delta \rightarrow 0} \sum_{j=1}^d \mathcal{E}_j \mathcal{T} \left(\mathcal{J} \frac{1}{\delta + \Gamma + \mathcal{L}_H - \mathcal{E} \cdot \nabla} (\nabla_j f_{\beta,\mu}(H)) \right)$$

This contains all non-linear terms in the electric field

Limit $\delta \rightarrow 0$ can be taken, if inverse exists

Linear coefficients of $j_{\beta,\mu,\mathcal{E}}$ in \mathcal{E} give conductivity tensor

In **relaxation time approximation** (RTA) one replaces Γ by $\frac{1}{\tau} > 0$

Theorem 7.4 (Kubo formula in RTA [BES])

$$\sigma_{i,j}(\beta, \mu, \tau) = \mathcal{T} \left(\nabla_i H \frac{1}{\frac{1}{\tau} + \mathcal{L}_H} (\nabla_j f_{\beta,\mu}(H)) \right)$$

Hall conductance $i \neq j$ at zero temperature $\beta = \infty$ and $\tau = \infty$ exists

$$\sigma_{i,j}(\beta = \infty, \mu, \tau = \infty) = \mathcal{T} \left((\mathcal{L}_H)^{-1} (\nabla_i H) \nabla_j P \right)$$

where $P = \chi(H \leq \mu)$. As

$$\nabla_j P = P \nabla_j P (\mathbf{1} - P) + (\mathbf{1} - P) \nabla_j P P$$

and

$$(\mathcal{L}_H)^{-1} (P \nabla_j H (\mathbf{1} - P)) = -i P \nabla_j P (\mathbf{1} - P)$$

$$(\mathcal{L}_H)^{-1} ((\mathbf{1} - P) \nabla_j H P) = i (\mathbf{1} - P) \nabla_j P P$$

Hence

$$\sigma_{i,j}(\beta = \infty, \mu, \tau = \infty) = i \mathcal{T} (P [\nabla_i P, \nabla_j P]) = \frac{1}{2\pi} \text{Ch}_{\{i,j\}}(P)$$

R.h.s. is integer-valued in dimension $d = 2$ and $d = 3$ (3D QHE)

This result holds also in a mobility gap regime [BES]

Electric polarization

$t \in [0, 2\pi) \cong \mathbb{S}^1 \mapsto H(t)$ periodic gapped Hamiltonian (changes dyn.)

Change ΔP in polarization is integrated induced current density:

$$\Delta P = \int_0^{2\pi} dt \mathcal{T}(\rho(t) J(t)) \quad , \quad \rho(0) = P_0 = \chi(H \leq \mu)$$

with $J(t) = i[H(t), X]$. Algebraic reformulation:

$$\Delta P = \int_0^{2\pi} dt \mathcal{T}(\rho(t) [\partial_t \rho(t), [X, \rho(t)]])$$

However, $\rho(t)$ unknown. So adiabatic limit of slow time changes:

Theorem 7.5 (Kingsmith-Vanderbilt and [ST])

$t \in \mathbb{S}^1 \mapsto H(t)$ smooth with gap open for all t

With $\rho(0) = P_0(0)$ and $\varepsilon \partial_t \rho(t) = i[\rho(t), H(t)]$, for any $N \in \mathbb{N}$

$$\Delta P = i \int_0^{2\pi} dt \mathcal{T}(P_0(t) [\partial_t P_0(t), [X, P_0(t)]]) + \mathcal{O}(\varepsilon^N)$$

Now add time to algebra: $C(\mathbb{S}^1, \mathcal{A}_d)$ is like \mathcal{A}_{d+1}

0th component is time and $\nabla_0 = \partial_t$

Also trace on $C(\mathbb{S}^1, \mathcal{A}_d)$ is $\frac{1}{2\pi} \int_0^{2\pi} dt \mathcal{T}$

Korollar 7.6

Polarization of periodically driven system is topological:

$$\Delta P_j = 2\pi \text{Ch}_{\{0,j\}} + \mathcal{O}(\varepsilon^N)$$

For $d = 1, 2$ and $j = 1$, one hence has $\Delta P_1 \in 2\pi \mathbb{Z}$ up to $\mathcal{O}(\varepsilon^N)$

However, in $d = 3$ one does **not** have $\Delta P_j \in 2\pi \mathbb{Z}$, but due to generalized Streda formula, magneto-electric response satisfies

$$\alpha_{1,2,3} = \partial_{B_{2,3}} \Delta P_1 = 2\pi \text{Ch}_{\{0,1,2,3\}} \in 2\pi \mathbb{Z}$$

Similarly: IDOS on gaps satisfies gap labelling

Chiral polarization

Chiral Hamiltonian $H = -\sigma_3 H \sigma_3$, typically due to sub-lattice symmetry
chiral polarization = difference between two electric dipole moments

$$P_C = \mathbf{E} \operatorname{Tr} \langle 0 | P \sigma_3 X P | 0 \rangle = i \mathcal{T}(P \sigma_3 \nabla P)$$

due to $X|0\rangle = 0$. Let U be Fermi unitary of P

Proposition 7.7 ([PS])

$$P_{C,j} = -\frac{1}{2} \operatorname{Ch}_{\{j\}}(U) \quad , \quad j = 1, \dots, d$$

Proof. Expressing P in terms of U

$$P_C = \frac{i}{4} \mathcal{T} \left(\left(\begin{array}{cc} \mathbf{1} & U^* \\ -U & -\mathbf{1} \end{array} \right) \left(\begin{array}{cc} 0 & -\nabla U^* \\ -\nabla U & 0 \end{array} \right) \right) = \frac{i}{4} \mathcal{T}(-U^* \nabla U + U \nabla U^*)$$

Now use $U \nabla U^* = -(\nabla U) U^*$ and cyclicity □

8 Bulk-boundary correspondence and applications

Toeplitz extension $T(\mathcal{A}_d) = C^*(S_1^B, \dots, S_{d-1}^B, \widehat{S}_d^B, W_\omega)$

$$\begin{array}{ccccccc}
 & & \text{edge} & & \text{half-space} & & \text{bulk} \\
 0 & \rightarrow & \mathcal{E}_d & \rightarrow & T(\mathcal{A}_d) & \rightarrow & \mathcal{A}_d \rightarrow 0
 \end{array}$$

Moreover: $\mathcal{E}_d \cong \mathcal{A}_{d-1} \otimes \mathcal{K}(\ell^2(\mathbb{N}))$

$$\begin{array}{ccccc}
 K_0(\mathcal{A}_{d-1}) & \xrightarrow{i_*} & K_0(T(\mathcal{A}_d)) & \xrightarrow{\pi_*} & K_0(\mathcal{A}_d) \\
 \uparrow \text{Ind} & & & & \downarrow \text{Exp} \\
 K_1(\mathcal{A}_d) & \xleftarrow{\pi_*} & K_1(T(\mathcal{A}_d)) & \xleftarrow{i_*} & K_1(\mathcal{A}_{d-1})
 \end{array}$$

Theorem 8.1 ([KRS, PS])

$$\text{Ch}_{I \cup \{d\}}(\mathbf{A}) = \text{Ch}_I(\text{Ind}(\mathbf{A})) \quad |I| \text{ even}, [\mathbf{A}] \in K_1(\mathcal{A}_d)$$

$$\text{Ch}_{I \cup \{d\}}(\mathbf{P}) = \text{Ch}_I(\text{Exp}(\mathbf{P})) \quad |I| \text{ odd}, [\mathbf{P}] \in K_0(\mathcal{A}_d)$$

Proof: loooong **Example:** $d = 1$ was exactly the SSH model

Physical implication in $d = 2$: QHE

P Fermi projection below a bulk gap $\Delta \subset \mathbb{R}$. Kubo formula:

$$\text{Hall conductance} = \text{Ch}_{\{1,2\}}(P)$$

Bulk-boundary:

$$\text{Ch}_{\{1,2\}}(P) = \text{Ch}_{\{1\}}(\text{Exp}(P)) = \text{Wind}(\text{Exp}(P))$$

With continuous $g(E) = 1$ for $E < \Delta$ and $g(E) = 0$ for $E > \Delta$:

$$\text{Exp}(P) = \exp(-2\pi i g(\hat{H})) \in T(\mathcal{A}_2)$$

as indeed $\pi(g(\hat{H})) = g(H) = P$ so that $\pi(\text{Exp}(P)) = \mathbf{1}$ trivial

Theorem 8.2 (Quantization of boundary currents [KRS, PS])

$$\text{Ch}_{\{1,2\}}(P) = \mathbb{E} \sum_{n_2 \geq 0} \langle 0, n_2 | g'(\hat{H}) i [X_1, \hat{H}] | 0, n_2 \rangle$$

The r.h.s. is current density flowing along the boundary

Proof: With $\widehat{\mathcal{T}}(A) = \mathcal{T}_1 \text{Tr}_2(A) = \mathbf{E}_{\mathbb{P}} \sum_{n_2 \geq 0} \langle 0, n_2 | \widehat{A}_\omega | 0, n_2 \rangle$, r.h.s. is

$$j^e(g) = \mathbb{E} \sum_{n_2 \geq 0} \langle 0, n_2 | g'(\widehat{H}) i[X_1, \widehat{H}] | 0, n_2 \rangle = \widehat{\mathcal{T}}(\widehat{J}_1 g'(\widehat{H}))$$

Summability in n_2 has to be checked

Let $\Pi : \ell^2(\mathbb{Z}^2) \rightarrow \ell^2(\mathbb{Z} \times \mathbb{N})$ surjective partial isometry,
namely $\Pi \Pi^*$ identity on $\ell^2(\mathbb{Z} \times \mathbb{N})$

Then $\widehat{H} = \Pi H \Pi^*$

Proposition 8.3

For $G \in C^\infty(\mathbb{R})$ with $\text{supp}(G) \cap \sigma(H) = \emptyset$

Then the operator $G(\widehat{H})$ is $\widehat{\mathcal{T}}$ -traceclass

Proof based on functional calculus often attributed to Helffer-Sjorstrand

Proposition 8.4 (Functional calculus à la Dynkin 1972)

$\chi \in C_0^\infty((-1, 1), [0, 1])$ even and equal to 1 on $[-\delta, \delta]$

For $N \geq 1$ let quasi-analytic extension $\tilde{G} : \mathbb{C} \rightarrow \mathbb{C}$ of G by

$$\tilde{G}(x, y) = \sum_{n=0, \dots, N} G^{(n)}(x) \frac{(iy)^n}{n!} \chi(y), \quad z = x + iy$$

Then with norm-convergent Riemann sum

$$G(H) = \frac{-1}{2\pi} \int_{\mathbb{R}^2} dx dy \partial_{\bar{z}} \tilde{G}(x, y) (z - H)^{-1}$$

Proof. Crucial identity is

$$\partial_{\bar{z}} \tilde{G}(x, y) = G^{(N+1)}(x) \frac{(iy)^N}{N!} \chi(y) + i \sum_{n=0, \dots, N} G^{(n)}(x) \frac{(iy)^n}{n!} \chi'(y)$$

In particular, uniformly in x, y , one has $|\partial_{\bar{z}} \tilde{G}(x, y)| \leq C |y|^N$

Hence also $\partial_{\bar{z}} \tilde{G}(x, 0) = 0$. Now resolvent bound. Details.... □

Proof of Proposition 8.3. Geometric resolvent identity

$$\frac{1}{z - \hat{H}} = \Pi \frac{1}{z - H} \Pi^* + \frac{1}{z - \hat{H}} (\hat{H} \Pi^* - \Pi H) \frac{1}{z - H} \Pi^*$$

in Dykin for $G(\hat{H})$ together with $G(H) = 0$ leads to

$$\begin{aligned} G(\hat{H}) &= \Pi G(H) \Pi^* + \hat{K} \\ &= \frac{-1}{2\pi} \int_{\mathbb{R}^2} dx dy \partial_{\bar{z}} \tilde{G}(x, y) \frac{1}{z - \hat{H}} (\hat{H} \Pi^* - \Pi H) \frac{1}{z - H} \Pi^* \end{aligned}$$

Resolvents have fall-off of their matrix elements off the diagonal:

$$(n_j - m_j)^k \langle n | (z - H)^{-1} | m \rangle = i^k \langle n | \nabla_j^k (z - H)^{-1} | m \rangle, \quad k \in \mathbb{N}$$

Expand $\nabla^k (z - H)^{-1}$ by Leibniz rule. As $\|\nabla^k H\| \leq C$

$$|\langle n | (z - H)^{-1} | m \rangle| \leq \frac{1}{|y|^{k+1}} \frac{C_k}{1 + |n_j - m_j|^k}$$

Same bound holds for resolvent of \hat{H} (improvement: Combes-Thomas)

If finite range, $\widehat{H}\Pi^* - \Pi H$ has matrix elements only on boundary. Then

$$\begin{aligned}
 & |\langle 0, n_2 | \widehat{K} | 0, n_2 \rangle| \\
 & \leq \sum_{m \in \mathbb{Z} \times \mathbb{N}} \sum_{k \in \mathbb{Z}^2} \frac{1}{2\pi} \int_{\mathbb{R}^2} dx dy |\partial_{\bar{z}} \tilde{G}(x, y)| |\langle 0, n_2 | (z - H)^{-1} | m \rangle| \\
 & \qquad \qquad \qquad |\langle m | \widehat{H}\Pi^* - \Pi H | k \rangle| |\langle k | (z - H)^{-1} | 0, n_2 \rangle| \\
 & \leq C \sum_{m_1 \geq 0} \int_{\mathbb{R}^2} dx dy |\partial_{\bar{z}} \tilde{G}(x, y)| \frac{1}{|y|^{2k+2}} \frac{1}{1 + |n_2|^{2k}} \frac{1}{1 + |m_1|^{2k}}
 \end{aligned}$$

Now above bound on resolvent for $N \geq 2k + 2$

As integral over bounded region, sum can be carried out

$$|\langle 0, n_2 | \widehat{K} | 0, n_2 \rangle| \leq \frac{C}{1 + |n_2|^{2k}}$$

But this implies desired $\widehat{\mathcal{T}}$ -traceclass estimate □

Proof of Theorem 8.2. Set $\hat{U} = \text{Exp}(P) = \exp(-2\pi i g(\hat{H}))$ and

$$\text{Ind} = i \hat{\mathcal{T}}((\hat{U}^* - \mathbf{1}) \nabla_1 \hat{U})$$

Express \hat{U} as exponential series and use Leibniz rule:

$$\text{Ind} = \sum_{m=0}^{\infty} \frac{(2\pi i)^m}{m!} \sum_{l=0}^{m-1} \hat{\mathcal{T}} \left((\hat{U}^* - \mathbf{1}) g(\hat{H})^l \nabla_1 g(\hat{H}) g(\hat{H})^{m-l-1} \right)$$

where trace and sum exchange by $\hat{\mathcal{T}}$ -traceclass property of $\hat{U} - \mathbf{1}$

Due to cyclicity and $[\hat{U}, g(\hat{H})] = 0$, each summand equal to

$$\hat{\mathcal{T}}((\hat{U}^* - \mathbf{1}) g(\hat{H})^{m-1} \nabla_1 g(\hat{H}))$$

Exchanging sum and trace, summing up again:

$$\text{Ind} = -2\pi \hat{\mathcal{T}} \left((\mathbf{1} - \hat{U}) \nabla_1 g(\hat{H}) \right)$$

Now same argument for $\hat{U}^k = \exp(-2\pi i k g(\hat{H}))$ for $k \neq 0$,

$$\text{Ind} = \frac{i}{k} \hat{\mathcal{T}}((\hat{U}^k - \mathbf{1})^* \nabla_1 \hat{U}^k) = -2\pi \hat{\mathcal{T}} \left((\mathbf{1} - \hat{U}^k) \nabla_1 g(\hat{H}) \right)$$

Writing $g(E) = \int dt \tilde{g}(t) e^{-E(1+it)}$ with adequate \tilde{g} , by DuHamel

$$\text{Ind} = 2\pi \int dt \tilde{g}(t) (1+it) \int_0^1 dq \hat{T} \left((\hat{U}^k - \mathbf{1}) e^{-(1-q)(1+it)\hat{H}} (\nabla_1 \hat{H}) e^{-q(1+it)\hat{H}} \right)$$

With $g'(E) = - \int dt (1+it) \tilde{g}(t) e^{-E(1+it)}$ for $k \neq 0$,

$$\text{Ind} = 2\pi \hat{T} \left((\hat{U}^k - \mathbf{1}) g'(\hat{H}) \nabla_1 \hat{H} \right)$$

For $k = 0$, the r.h.s. vanishes. To conclude, let $\phi \in C_0^\infty((0, 1), \mathbb{R})$

Fourier coefficients $a_k = \int_0^1 dx e^{-2\pi ikx} \phi(x)$ satisfy $\sum_k a_k e^{2\pi ikx} = \phi(x)$

In particular, $\sum_k a_k = 0$ and

$$\begin{aligned} a_0 \text{Ind} &= - \sum_{k \neq 0} a_k \text{Ind} = 2\pi \sum_k a_k \hat{T} \left((\mathbf{1} - \hat{U}^k) g'(\hat{H}) \nabla_1 \hat{H} \right) \\ &= 2\pi \hat{T} \left((0 - \phi(g(\hat{H}))) g'(\hat{H}) \nabla_1 \hat{H} \right) \end{aligned}$$

As $\phi \rightarrow \chi_{[0,1]}$ also $a_0 \rightarrow 1$ and $\phi(g(\hat{H}))g'(\hat{H}) \rightarrow g'(\hat{H})$ (no Gibbs)

As $J_1 = \nabla_1 \hat{H}$ proof is concluded □

Chiral system in $d = 3$: anomalous surface QHE

Chiral Fermi projection P (off-diagonal) \implies Fermi unitary A

$$\text{Ch}_{\{1,2,3\}}(A) = \text{Ch}_{\{1,2\}}(\text{Ind}(A))$$

Magnetic field perpendicular to surface opens gap in surface spec.

With $\hat{P} = \hat{P}_+ + \hat{P}_-$ projection on central surface band, as in SSH:

$$\text{Ind}(A) = [\hat{P}_+] - [\hat{P}_-]$$

Theorem 8.5 ([PS])

Suppose either $\hat{P}_+ = 0$ or $\hat{P}_- = 0$ (conjectured to hold). Then:

$\text{Ch}_{\{1,2,3\}}(A) \neq 0 \implies$ surface QHE, Hall cond. imposed by bulk

Actually only approximate chiral symmetry needed

Experiment? No (approximate) chiral topological material known

Delocalization of boundary states

Hypothesis: bulk gap at Fermi level μ

Disorder: in arbitrary finite strip along boundary hypersurface

Theorem 8.6 ([PS])

For even d , if strong invariant $\text{Ch}_{\{1, \dots, d\}}(P) \neq 0$,

then no Anderson localization of boundary states in bulk gap

Technically: Aizenman-Molcanov bound for no energy in bulk gap

Theorem 8.7 ([PS])

For odd $d \geq 3$, if strong invariant $\text{Ch}_{\{1, \dots, d\}}(A) \neq 0$,

then no Anderson localization at $\mu = 0$

BBC for periodically driven systems

In time direction: stroboscopes Here: in spacial direction

Lift $t \in \mathbb{S}^1 \cong [0, 2\pi) \mapsto \hat{H}(t)$ of $t \in \mathbb{S}^1 \mapsto H(t)$ in

$$0 \longrightarrow C(\mathbb{S}^1, \mathcal{E}_d) \xrightarrow{i} C(\mathbb{S}^1, \hat{\mathcal{A}}_d) \xrightarrow{\text{ev}} C(\mathbb{S}^1, \mathcal{A}_d) \longrightarrow 0$$

Then for polarization in direction d with adiabatic projection P_A :

$$\Delta P_d = 2\pi \text{Ch}_{\{0,d\}}(P_A) = 2\pi \text{Ch}_{\{0\}}(U_\Delta)$$

where 0-th component still time and $[U_\Delta]_1 = \text{Exp}[P_A]_0$. Now

$$\text{Ch}_{\{0\}}(U_\Delta) = -2\pi \int_0^{2\pi} dt \hat{T} \left(g'(\hat{H}(t)) \partial_t \hat{H}(t) \right)$$

For $d = 1$, this is 2π times spectral flow of boundary eigenvalues. Thus

$$\Delta P_1 = -2\pi \text{SF}(t \in \mathbb{S}^1 \mapsto \hat{H}(t) \text{ by } \mu)$$

namely charge pumped from valence to conduction states

For $d > 1$, spectral flow is in sense of Breuer-Fredholm operators

9 Implementation of symmetries

This invokes real structure simply denoted by bar on \mathcal{H} and $\mathcal{B}(\mathcal{H})$

$$\text{chiral symmetry (CHS)} : \quad J_{\text{ch}}^* H J_{\text{ch}} = -H$$

$$\text{time reversal symmetry (TRS)} : \quad S_{\text{tr}}^* \bar{H} S_{\text{tr}} = H$$

$$\text{particle-hole symmetry (PHS)} : \quad S_{\text{ph}}^* \bar{H} S_{\text{ph}} = -H$$

$S_{\text{tr}} = e^{i\pi s^y}$ orthogonal on \mathbb{C}^{2s+1} with $S_{\text{tr}}^2 = \pm 1$ even or odd

S_{ph} orthogonal on \mathbb{C}_{ph}^2 with $S_{\text{ph}}^2 = \pm 1$ even or odd

Note: TRS + PHS \implies CHS with $J_{\text{ch}} = S_{\text{tr}} S_{\text{ph}}$

10 combinations of symmetries: none (1), one (5), three (4)

10 Cartan-Altland-Zirnbauer classes (CAZ): 2 complex, 8 real

Further distinction in each of the 10 classes: topological insulators

Periodic table of topological insulators

Schnyder-Ryu-Furusaki-Ludwig, Kitaev 2008: just strong invariants

$j \setminus d$	TRS	PHS	CHS	1	2	3	4	5	6	7	8
0	0	0	0		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}
1	0	0	1	\mathbb{Z}		\mathbb{Z}		\mathbb{Z}		\mathbb{Z}	
0	+1	0	0				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}
1	+1	+1	1	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2
2	0	+1	0	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2
3	-1	+1	1	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$	
4	-1	0	0		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$
5	-1	-1	1	$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}			
6	0	-1	0		$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}		
7	+1	-1	1			$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}	

Periodic table: real classes only

64 pairings = 8 KR-cycles paired with 8 KR-groups

$j \setminus d$	TRS	PHS	CHS	1	2	3	4	5	6	7	8
0	+1	0	0				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}
1	+1	+1	1	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2
2	0	+1	0	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$		\mathbb{Z}_2
3	-1	+1	1	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$	
4	-1	0	0		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}				$2\mathbb{Z}$
5	-1	-1	1	$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}			
6	0	-1	0		$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}		
7	+1	-1	1			$2\mathbb{Z}$		\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}	

Focus on system in $d = 2$ with odd TRS $S = S_{\text{tr}}$:

$$S^2 = -1 \quad S^* \bar{H} S = H$$

\mathbb{Z}_2 index for odd TRS and $d = 2$

Rewrite $S^* \bar{H} S = H = S^* H^t S$ with $H^t = (\bar{H})^*$

$\implies S^* (H^n)^t S = H^n$ for $n \in \mathbb{N} \implies S^* P^t S = P$

For $d = 2$, Dirac phase $F = \frac{X_1 + iX_2}{|X_1 + iX_2|} = F^t$ and $[S, F] = 0$

Hence Fredholm operator $T = PFP$ of following type

Definition T odd symmetric $\iff S^* T^t S = T \iff (TS)^t = -TS$

Theorem 9.1 (Atiyah-Singer 1969)

$\mathbb{F}_2(\mathcal{H}) = \{\text{odd symmetric Fredholm operators}\}$ has 2 connected components labelled by compactly stable homotopy invariant

$$\text{Ind}_2(T) = \dim(\text{Ker}(T)) \bmod 2 \in \mathbb{Z}_2$$

Application: \mathbb{Z}_2 phase label for Kane-Mele model if dyn. localized

Existence proof of \mathbb{Z}_2 -indices via Kramers arg.

First of all: $\text{Ind}(T) = 0$ because $\text{Ker}(T^*) = S \overline{\text{Ker}(T)}$

Idea: $\text{Ker}(T) = \text{Ker}(T^* T)$

and positive eigenvalues of $T^* T$ have even multiplicity

Let $T^* T v = \lambda v$ and $w = S \overline{T v}$ (N.B. $\lambda \neq 0$). Then

$$\begin{aligned} T^* T w &= S (S^* T^* S) (S^* T S) \overline{T v} \\ &= S \overline{T T^* T v} = \lambda S \overline{T v} = \lambda w. \end{aligned}$$

Suppose now $\mu \in \mathbb{C}$ with $v = \mu w$. Then

$$v = \mu S \overline{T v} = \mu S \overline{T \overline{\mu} S T v} = -|\mu|^2 T^* T v = -|\mu|^2 \lambda v$$

Contradiction to $v \neq 0$.

Now $\text{span}\{v, w\}$ is invariant subspace of $T^* T$.

Go on to orthogonal complement

Symmetries of the Dirac operator

$$D = \sum_{j=1}^d X_j \otimes \mathbf{1} \otimes \gamma_j$$

$\gamma_1, \dots, \gamma_d$ irrep of C_d with $\gamma_{2j} = -\overline{\gamma_{2j}}$ and $\gamma_{2j+1} = \overline{\gamma_{2j+1}}$

In even d exists grading $\Gamma = \Gamma^*$ with $D = -\Gamma D \Gamma$ and $\Gamma^2 = \mathbf{1}$

Moreover, exists real unitary Σ (essentially unique) with

$d = 8 - i$	8	7	6	5	4	3	2	1
Σ^2	1	1	-1	-1	-1	-1	1	1
$\Sigma^* \overline{D} \Sigma$	D	$-D$	D	D	D	$-D$	D	D
$\Gamma \Sigma \Gamma$	Σ		$-\Sigma$		Σ		$-\Sigma$	

(D, Γ, Σ) defines a KR^i -cycle (spectral triple with real structure)

(Kasparov 1981, Connes 1995, Gracia-Varilly-Figueroa 2000)

Index theorems for periodic table

Symmetries of KR -cycles **and** Fermi projection/unitary lead to:

Theorem 9.2

Index theorems for all strong invariants in periodic table

Remarks:

Result holds also in the regime of strong Anderson localization
 $2\mathbb{Z}$ entries result from quaternionic Fredholm (even Ker, CoKer)

Links to Atiyah-Singer classifying spaces

Formulation as Clifford valued index theorem possible

Physical implications: case by case study necessary!

Example: focus on TRS $d = 2$ quantum spin Hall system (QSH)

Spin Chern numbers [Pro]

Approximate spin conservation \implies spin Chern numbers $\text{SCh}(P)$

Kane-Mele Hamiltonian has small commutator $[H, s_z]$

Also $[P, s_z]$ small and thus $Ps_zP|_{\text{Ran}(P)}$ spectrum close to $\{-1, 1\}$

\implies spectral gap! Let P_{\pm} be two associated spectral projections

Proposition 9.3 ([Pro])

P_{\pm} have off-diagonal decay so that Chern numbers can be defined

Hence $P = P_+ + P_-$ decomposes in two *smooth* projections

Definition 9.4

Spin Chern number of P is $\text{SCh}(P) = \text{Ch}(P_+)$

By TRS, $\text{Ch}(P) = 0$ and thus $\text{SCh}(P) = -\text{Ch}(P_-)$

Theorem 9.5 ([SB3])

$\text{Ind}_2(PFP) = \text{SCh}(P) \bmod 2$

Spin filtered helical edge channels for QSH

Remarkable: Non-trivial topology $SCh(P)$ persists TRS breaking!

General strategy: approximately conserved quantities lead to integer-valued invariants which persist breaking of real symmetry

Further example:

Kitaev chain (Class D with \mathbb{Z}_2 -invariant) has a winding number

Theorem 9.6

If $SCh(P) \neq 0$, spin filtered edge currents in $\Delta \subset \text{gap}$ are stable w.r.t. perturbations by magnetic field and disorder:

$$\mathbf{E} \text{Tr} \langle 0 | \chi_{\Delta}(\hat{H}) \frac{1}{2} \{i[\hat{H}, X_1], s_z\} | 0 \rangle = |\Delta| SCh(P) + \text{correct.}$$

Resumé: $\text{Ind}_2(PFP) = 1 \implies$ no Anderson loc. for edge states

Rice group of Du (since 2011): QSH stable w.r.t. magnetic field

10 Laughlin arguments


Satz 10.1 ([DS])

H disordered Harper-like operator on $\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^L$ with $\mu \in \text{gap}$


H_α Hamiltonian with extra flux $\alpha \in [0, 1]$ through 1 cell of \mathbb{Z}^2

Then for $P = \chi(H \leq \mu)$

$$\text{SF}(\alpha \in [0, 1] \mapsto H_\alpha \text{ through } \mu) = -\text{Ch}_{\{1,2\}}(P)$$



Phillips' analytic definition (1996)



\exists finite partition $0 = t_0 < t_1 < \dots < t_{N-1} < t_N = 1$ of $[0, 1]$ and $a_n < 0 < b_n$ with $t \in [t_{n-1}, t_n] \mapsto \chi(T_t \in [a_n, b_n])$ continuous. Set:

$$\text{SF}(t \in [0, 1] \mapsto T_t) = \sum_{n=1}^N \text{Tr}_{\mathcal{H}} (\chi(T_{t_{n-1}} \in [a_n, 0]) - \chi(T_{t_n} \in [a_n, 0]))$$

Theorem 10.2 (Phillips 1996)

$SF(t \in [0, 1] \mapsto T_t)$ independent of partition and $a_n < 0 < b_n$.

It is a homotopy invariant when end points are kept fixed.

It satisfies concatenation and normalization:

$$SF(t \in [0, 1] \mapsto T + (1 - 2t)P) = -\dim(P) \quad \text{for } TP = P$$

Theorem 10.3 (Lesch 2004)

Homotopy invariance, concatenation, normalization characterize SF

Theorem 10.4 (Perera 1993, Phillips 1996)

SF on loops establishes isomorphism $\pi_1(\mathbb{F}_{sa}^) = \mathbb{Z}$*

Theorem 10.5 (Avron-Seiler-Simon 1994, Phillips 1996)

0 gap of $H = H^*$ and $P = \chi(H \leq 0)$. If $t \in [0, 1] \mapsto H_t$ with

- (i) $H_1 = UH_0U^*$ for unitary U
- (ii) 0 in essential gap of H_t for all $t \in [0, 1]$

then

$$\text{SF}\left(t \in [0, 1] \mapsto H_t \text{ through } 0\right) = -\text{Ind}(PUP)$$

Exact sequence interpretation: Mapping cone associated to U :

$$\mathcal{M} = \{t \in [0, 1] \mapsto A_t \in \mathcal{A} + \mathcal{K} : A_0 = U^*A_1U, A_t - A_0 \in \mathcal{K}\}$$

with $0 \rightarrow SK \hookrightarrow \mathcal{M} \xrightarrow{\text{ev}} \mathcal{A} \rightarrow 0$. Now $K_1(SK) = K_0(\mathcal{K}) = \mathbb{Z}$ and


$$\text{Exp}[P]_0 = [\exp(2\pi i \text{Lift}(P)_t)]_1 = [\exp(2\pi i(P + tU^*[P, U]))]_1$$

Then

$$\langle (\mathcal{H}, F), [P]_0 \rangle = \left\langle \left(\int dt \otimes \text{Tr}, \partial_t \right), \text{Exp}[P]_0 \right\rangle = \text{SF}(2P-1+tU^*[2P-1, U])$$

Proof of bulk-boundary in $d = 2$ (idea Macris 2002)

Based on gauge invariance and compact stability




\mathbb{Z}_2 invariant and \mathbb{Z}_2 spectral flow for QSH

Theorem 10.6


$\alpha \in [0, 1] \mapsto H(\alpha)$ inserted flux in Kane-Mele model (breaks TRS)

$\text{Ind}_2(\text{PFP}) = 1 \implies H(\alpha = \frac{1}{2})$ has TRS + Kramers pair in gap



Higher dimensional Laughlin arguments

$d = 1$: chiral spectral flow in SSH leads to bound state of $H_{\frac{1}{2}}$



On $\ell^2(\mathbb{Z}^d) \otimes \mathbb{C}^L$ with $d \geq 3$: insert non-abelian Wu-Yang monopol

$$A = \frac{i}{2} \frac{[D, \gamma]}{D^2}, \quad D = \sum_{j=1}^d \gamma_j X_j$$

into non-abelian translations (say without magnetic field):

$$S_k^\alpha = e^{i\nabla_k^\alpha} = U^\alpha(X) S_k, \quad \nabla_k^\alpha = i\partial_k + \alpha A_k$$

Then study (chiral) spectral flow for $H_\alpha = P(S_1^\alpha, \dots, S_d^\alpha)$

11 Dirty superconductors

Disordered one-electron Hamiltonian h on $\mathcal{H} = \ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^{2s+1}$

$c = (c_{n,l})$ annihilation operators on fermionic Fock space $\mathcal{F}_-(\mathcal{H})$

Hamilt. on $\mathcal{F}_-(\mathcal{H})$ with mean field pair creation $\Delta^* = -\bar{\Delta} \in \mathcal{B}(\mathcal{H})$

$$\begin{aligned} \mathbf{H} - \mu \mathbf{N} &= c^* (h - \mu \mathbf{1}) c + \frac{1}{2} c^* \Delta c^* - \frac{1}{2} c \bar{\Delta} c \\ &= \frac{1}{2} \begin{pmatrix} c \\ c^* \end{pmatrix}^* \begin{pmatrix} h - \mu & \Delta \\ -\bar{\Delta} & -\bar{h} + \mu \end{pmatrix} \begin{pmatrix} c \\ c^* \end{pmatrix} \end{aligned}$$

Hence BdG Hamiltonian on $\mathcal{H}_{\text{ph}} = \mathcal{H} \otimes \mathbb{C}_{\text{ph}}^2$

$$H_{\mu} = \begin{pmatrix} h - \mu & \Delta \\ -\bar{\Delta} & -\bar{h} + \mu \end{pmatrix}$$

Even PHS (Class D)

$$S_{\text{ph}}^* \bar{H}_{\mu} S_{\text{ph}} = -H_{\mu} \quad , \quad S_{\text{ph}} = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$$

Class D systems

$\text{spec}(H_\mu) = -\text{spec}(H_\mu)$ and generically gap or pseudo-gap at 0

Satz 11.1

Gibbs (KMS) state for observable $\mathbf{Q} = d\Gamma(Q)$

$$\frac{1}{Z_{\beta,\mu}} \text{Tr}_{\mathcal{F}_-(\mathcal{H})} \left(\mathbf{Q} e^{-\beta(\mathbf{H}-\mu\mathbf{N})} \right) = \text{Tr}_{\mathcal{H}_{\text{ph}}} (f_\beta(H_\mu) Q)$$

Example $p + ip$ wave superconductor with $\mathcal{H} = \ell^2(\mathbb{Z}^2)$

$$h = \mathbf{S}_1 + \mathbf{S}_1^* + \mathbf{S}_2 + \mathbf{S}_2^* \quad \Delta_{p+ip} = \delta (\mathbf{S}_1 - \mathbf{S}_1^* + i(\mathbf{S}_2 - \mathbf{S}_2^*))$$

Then $P = \chi(H_\mu \leq 0)$ satisfies $\text{Ch}(P) = 1$ for $\mu > 0$ and $\delta > 0$

Conjecture (Kubo missing) Quantized Wiedemann-Franz

$$\kappa_H = \frac{\pi}{8} \text{Ch}(P) T + \mathcal{O}(T^2)$$

Spectral flow in a BdG-Hamiltonian

Flux tube in two-dimensional BdG Hamiltonian

$$S_{\text{ph}}^* \overline{H_\alpha} S_{\text{ph}} = -H_{-\alpha} \quad , \quad S_{\text{ph}}^2 = \pm 1$$

Then $S_{\text{ph}}^* \overline{H_\alpha} S_{\text{ph}} = -U^* H_{1-\alpha} U$ so that

$$\sigma(H_\alpha) = -\sigma(H_{-\alpha}) = -\sigma(H_{1-\alpha})$$

PHS only for $\alpha = 0, \frac{1}{2}, 1$ and thus $\text{Ind}_2(H_{\frac{1}{2}})$ well-defined

Theorem 11.2 ([DS])

$$\text{Ind}(PUP) \bmod 2 = \text{Ind}_2(H_{\frac{1}{2}})$$

or: odd Chern number implies existence of zero mode at defect

These zero modes are Majorana fermions (Read-Green 2000)

Worth noting: $S_{\text{ph}}^2 = -1 \implies \text{Ind}(PUP) \text{ even} \implies \text{no zero mode}$

Spin quantum Hall effect in Class C

Satz 11.3 (Altland-Zirnbauer 1997)

$SU(2)$ spin rotation invariance $[\mathbf{H}, \mathbf{s}] = 0$

$\implies H = H_{\text{red}} \otimes \mathbf{1}$ with odd PHS (Class C)

$$\mathcal{S}_{\text{ph}}^* \overline{H_{\text{red}}} \mathcal{S}_{\text{ph}} = -H_{\text{red}} \quad , \quad \mathcal{S}_{\text{ph}} = \begin{pmatrix} 0 & -\mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix}$$

Example $d + id$ wave superconductor with h as above and

$$\Delta_{d+id} = \delta (i(\mathcal{S}_1 + \mathcal{S}_1^* - \mathcal{S}_2 - \mathcal{S}_2^*) + (\mathcal{S}_1 - \mathcal{S}_1^*)(\mathcal{S}_2 - \mathcal{S}_2^*)) s^2$$

Again $\text{Ch}(P) = 2$ for $\delta > 0$ and $\mu > 0$

Satz 11.4

Spin Hall conductance (Kubo) and spin edge currents quantized

Current aims:

- analysis of topology associated to spacial reflections, etc.
- Index theory for weak invariants via KK -theory
- bulk-edge correspondence in real cases
- further investigation of physical implications of invariants
- stability of invariants w.r.t. interactions
- analysis of bosonic systems and photonic crystals

Physics References

- [KM] C. L. Kane, E. J. Mele, *Quantum spin Hall effect in graphene*, Phys. Rev. Lett. **95**, 226801 (2005), *Z(2) topological order and the quantum spin Hall effect*, Phys. Rev. Lett. **95**, 146802 (2005).
- [RSFL] S. Ryu, A. P. Schnyder, A. Furusaki, A. W. W. Ludwig, *Topological insulators and superconductors: tenfold way and dimensional hierarchy*, New J. Phys. **12**, 065010 (2010).
- [Kit] A. Kitaev, *Periodic table for topological insulators and superconductors*, (Advances in Theoretical Physics: Landau Memorial Conference) AIP Conf. Proc. **1134**, 22-30 (2009).
- [SSH] W. P. Su, J. R. Schrieffer, A. J. Heeger, *Soliton excitations in polyacetylene*, Phys. Rev. **B 22**, 2099-2111 (1980).
- [AZ] A. Altland and M. R. Zirnbauer, *Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures*, Phys. Rev. B **55**, 1142-1161 (1997).

General Mathematics References

- [BR] O. Bratteli, D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1*, (Springer, Berlin, 1979).
- [CP] A. L. Carey, J. Phillips, *Spectral flow in Fredholm modules, eta invariants and the JLO cocycle*, *K-Theory* **31**, 135-194 (2004).
- [Con] A. Connes, *Noncommutative Geometry*, (Academic Press, San Diego, 1994).
- [GVF] J. M. Gracia-Bondía, J. C. Várilly, H. Figueroa, *Elements of noncommutative geometry*, (Springer Science & Business Media, 2013).
- [RLL] M. Rordam, F. Larsen, N. Laustsen, *An Introduction to K-theory for C^* -algebras*, (Cambridge University Press, Cambridge, 2000).
- [WO] N. E. Wegge-Olsen, *K-theory and C^* -algebras*, (Oxford Univ. Press, Oxford, 1993).

References Schulz-Baldes *et. al.*

- [PS] E. Prodan, H. Schulz-Baldes, *Bulk and boundary invariants for complex topological insulators: From K-theory to physics*, (Springer Int. Pub., Switzerland, 2016).
- [BES] J. Bellissard, A. van Elst, H. Schulz-Baldes, *The non-commutative geometry of the quantum Hall effect*, J. Math. Phys. **35**, 5373-5451 (1994).
- [KRS] J. Kellendonk, T. Richter, H. Schulz-Baldes, *Edge current channels and Chern numbers in the integer quantum Hall effect*, Rev. Math. Phys. **14**, 87-119 (2002).
- [LSB] T. Loring, H. Schulz-Baldes, *Finite volume calculation of K-theory invariants*, arXiv 2017.
- [GS] J. Grossmann, H. Schulz-Baldes, *Index pairings in presence of symmetries with applications to topological insulators*, Commun. Math. Phys. **343**, 477-513 (2016).

References Schulz-Baldes *et. al.*

- [SB1] H. Schulz-Baldes, *Topological insulators from the perspective of non-commutative geometry and index theory*, Jahresber Dtsch Math-Ver **118**, 247273 (2016)
- [SB2] H. Schulz-Baldes, *Persistence of spin edge currents in disordered quantum spin Hall systems*, Commun. Math. Phys. **324**, 589-600 (2013).
- [ST] H. Schulz-Baldes, S. Teufel, *Orbital polarization and magnetization for independent particles in disordered media*, Commun. Math. Phys. **319**, 649-681 (2013).
- [DS] G. De Nittis, H. Schulz-Baldes, *Spectral flows associated to flux tubes*, Annales H. Poincare **17**, 1-35 (2016).
- [CPS] A. L. Carey, J. Phillips, H. Schulz-Baldes, *Spectral flow for real skew-adjoint Fredholm operators*, J. Spec. Theory, to appear.
- [SB3] H. Schulz-Baldes, *\mathbb{Z}_2 -indices of odd symmetric Fredholm operators*, Dokumenta Math. **20**, 1481-1500 (2015).

More Mathematical Physics References

- [Pro] E. Prodan, *Robustness of the spin-Chern number*, Phys. Rev. **B** **80**, 125327 (2009).
- [BCR] C. Bourne, A. L. Carey, A. Rennie, *A noncommutative framework for topological insulators*, Rev. Math. Phys. **28**, 1650004 (2016).
- [BKR] C. Bourne, J. Kellendonk, A. Rennie, *The K-Theoretic Bulk-Edge Correspondence for Topological Insulators*, Ann. Henri Poincaré **18**, 1-34 (2017).
- [Lor] T. A. Loring, *K-theory and pseudospectra for topological insulators*, Annals of Physics **356**, 383-416 (2015).

Other groups (each with personal point of view)

- Bourne, Carey, Rennie, Kellendonk
- Mathai, Thiang, Hanabus
- Zirnbauer, Kennedy
- Panati, Monaco, Teufel, Cornean
- Katsura, Koma
- Hayashi, Furuta, Kotani
- Graf, Porta
- Gawedzki *et. al.*
- Kaufmann's, Li

- many theoretical physics groups